

Interrelationships Between Orthogonal, Unitary and Symplectic Matrix Ensembles

PETER J. FORRESTER AND ERIC M. RAINS

ABSTRACT. We consider the following problem: When do alternate eigenvalues taken from a matrix ensemble themselves form a matrix ensemble? More precisely, we classify all weight functions for which alternate eigenvalues from the corresponding orthogonal ensemble form a symplectic ensemble, and similarly classify those weights for which alternate eigenvalues from a union of two orthogonal ensembles forms a unitary ensemble. Also considered are the k -point distributions for the decimated orthogonal ensembles.

1. Introduction

Given a probability measure on a space of matrices, the eigenvalue PDF (probability density function) follows by a change of variables. For example, consider the space of $n \times n$ real symmetric matrices $A = [a_{j,k}]_{0 \leq j,k < n}$ with probability measure proportional to

$$e^{-\text{Tr}(A^2)/2}(dA), \quad (dA) := \prod_{j \leq k} a_{jk}. \quad (1-1)$$

The eigenvalues $x_0 < \dots < x_{n-1}$ are introduced via the spectral decomposition $A = RLR^T$ where R is a real orthogonal matrix with columns given by the eigenvectors of A and $L = \text{diag}(x_0, \dots, x_{n-1})$. Since $e^{-\text{Tr}(A^2)/2} = e^{-\sum_{j=0}^{n-1} x_j^2/2}$ the change of variables is immediate for the weight function; however the change of variables in (dA) cannot be carried out with such expedience.

The essential point of the latter task is to compute the Jacobian for the change of variables from the independent elements of A to the eigenvalues and the independent variables associated with the eigenvectors. Also, because only the eigenvalue PDF is being computed, one must integrate out the eigenvector dependence. In fact the dependence in the Jacobian on the eigenvalues separates from the dependence on the eigenvectors, so the task of performing the

integration does not become an issue. Explicitly, one finds (see e.g. [21])

$$(dA) = \Delta(x) \prod_{j=0}^{n-1} dx_j (R^T dR)$$

where $\Delta(x) := \Delta(x_0, \dots, x_{n-1}) := \prod_{0 \leq j < k < n} (x_k - x_j)$, and thus the eigenvalue PDF corresponding to (1-1) is proportional to

$$\prod_{j=0}^{n-1} g(x_j) |\Delta(x)|, \quad (1-2)$$

with $g(x) = e^{-x^2/2}$ (taking the absolute value of $\Delta(x)$ allows the ordering restriction on the eigenvalues to be dropped).

More generally, the above working shows that a space of $n \times n$ real symmetric matrices with probability measure proportional to

$$\exp\left(\sum_{j=1}^{\infty} \alpha_j \text{Tr}(A^j)\right) (dA) \quad (1-3)$$

will have eigenvalue PDF (1-2) with $g(x) = \exp(\sum_{j=1}^{\infty} \alpha_j x^j)$. Because (1-3) is unchanged by similarity transformations $A \mapsto RAR^T$ with R real orthogonal, for general g (1-2) is said to be the eigenvalue PDF of an orthogonal ensemble, and denoted $\text{OE}_n(g)$.

Real symmetric matrices are Hermitian matrices with all elements constrained to be real. If one considers Hermitian matrices without this constraint, so the off diagonal elements can now be complex, the eigenvalue PDF corresponding to the ensemble (1-3) is again given by (1-2) but with $|\Delta(x)|$ replaced by $(\Delta(x))^2$. Because (1-3) is then unchanged by $A \mapsto UAU^\dagger$ for U unitary, (1-2) so modified is referred to as a unitary ensemble and denoted $\text{UE}_n(g)$. The third and final possibility [7] is to consider $n \times n$ Hermitian matrices in which each element is itself a 2×2 matrix of the form

$$\begin{bmatrix} z & w \\ -\bar{w} & \bar{z} \end{bmatrix}.$$

This class of 2×2 matrices form the real quaternion number field \mathbb{H} . The spectrum of such matrices, regarded as $2n \times 2n$ matrices with complex entries, is doubly degenerate. The ensemble of matrices (1-3) is now invariant under the transformations $A \mapsto BAB^\dagger$ for B symplectic unitary, and so referred to as a symplectic ensemble. The eigenvalue PDF of the distinct eigenvalues is given by (1-2) with $|\Delta(x)|$ replaced by $(\Delta(x))^4$, and this is denoted $\text{SE}_n(g)$.

The matrix ensembles corresponding to the eigenvalue PDFs

$$\text{OE}_n(e^{-x^2/2}), \quad \text{UE}_n(e^{-x^2}), \quad \text{SE}_n(e^{-x^2})$$

are given the special labels GOE_n , GUE_n and GSE_n respectively (the G standing for Gaussian). As seen from (1–1) they can be realized by an appropriate Gaussian weight function in the probability space. Because for A real symmetric

$$e^{-\text{Tr}(A^2)/2} = \prod_{j=0}^{n-1} e^{-a_{jj}^2/2} \prod_{j < k}^{n-1} e^{-a_{jk}^2},$$

and similarly for A Hermitian with complex or real quaternion elements, independent elements of the Gaussian ensemble are independently distributed Gaussian random variables.

There are also a number of other known random matrix ensembles with this latter property, and which have eigenvalue PDF of the form $\text{OE}_n(g)$, $\text{UE}_n(g)$ or $\text{SE}_n(g)$ for some g . Seven such ensembles result by taking the Hermitian part of the matrix Lie algebras related to Cartan’s ten families of infinite symmetric spaces [30]. We specify five of them:

- $\text{Mat}(p, q; \mathbb{R})$ $p \times q$ matrices over \mathbb{R} ($p \geq q$),
- $\text{Mat}(p, q; \mathbb{C})$ $p \times q$ matrices over \mathbb{C} ($p \geq q$),
- $\text{Mat}(p, q; \mathbb{H})$ $p \times q$ matrices over \mathbb{H} ($p \geq q$),
- $\text{Symm}(n; \mathbb{C})$ $n \times n$ symmetric complex matrices,
- $\text{Anti}(n; \mathbb{C})$ $n \times n$ antisymmetric complex matrices.

The quantities of interest are the square of the non-zero singular values, or equivalently the eigenvalues of $A^\dagger A$ for A a member of the ensemble, in each case. The first two of these ensembles were studied long ago in mathematical statistics [29; 14]; these two together with the third have occurred in recent physical applications (see [3] and references therein), while the final two (in a different guise) have also arisen in a physical context [3]. The distribution of the eigenvalues of $A^\dagger A$ can be computed in a number of ways; one approach is to make use of the correspondence [30] to a symmetric space (of types BDI , $AIII$, CII , CI and $DIII$ respectively), which allows the tables in [16] to be utilized. Abusing notation, we have

$$\begin{aligned} \text{Mat}(p, q; \mathbb{R}) &= \text{OE}_q(x^{(p-q-1)/2} e^{-x/2}), \\ \text{Mat}(p, q; \mathbb{C}) &= \text{UE}_q(x^{p-q} e^{-x}), \\ \text{Mat}(p, q; \mathbb{H}) &= \text{SE}_q(x^{2(p-q)+1} e^{-x}), \\ \text{Symm}(n; \mathbb{C}) &= \text{OE}_n(e^{-x/2}), \\ \text{Anti}(2n; \mathbb{C}) &= \text{SE}_n(e^{-x}), \\ \text{Anti}(2n + 1; \mathbb{C}) &= \text{SE}_n(x^2 e^{-x}). \end{aligned} \tag{1–4}$$

Up to the scale of x , all the above weight functions are of the Laguerre form $x^\alpha e^{-x}$ and so by definition are examples of Laguerre matrix ensembles.

Another class of matrix ensembles in which the entries of the underlying matrices are independently distributed Gaussian random variables are known in mathematical statistics [21]. With $a \in \text{Mat}(p_1, q; \mathbb{F})$ (where $\mathbb{F} = \mathbb{R}, \mathbb{C}$ or \mathbb{H}), $b \in \text{Mat}(p_2, q, \mathbb{F})$, and $A = a^\dagger a$, $B = b^\dagger b$, these distributions are described by

$$\text{Beta}(p_1, p_2, q; \mathbb{F}) \quad q \times q \text{ matrices } A(A + B)^{-1}.$$

They have corresponding eigenvalue PDF (abusing notation as in (1–4))

$$\begin{aligned} \text{Beta}(p_1, p_2, q; \mathbb{R}) &= \text{OE}_q(x^{(p_1-q-1)/2}(1-x)^{(p_2-q-1)/2}), \\ \text{Beta}(p_1, p_2, q; \mathbb{C}) &= \text{UE}_q(x^{p_1-q}(1-x)^{p_2-q}), \\ \text{Beta}(p_1, p_2, q; \mathbb{H}) &= \text{SE}_q(x^{2(p_1-q)+1}(1-x)^{2(p_2-q)+1}), \end{aligned}$$

where $0 < x < 1$, and thus involve weight functions of the Jacobi type.

The revision above demonstrates that it is possible to realize, in terms of matrices with entries which are independently distributed Gaussian random variables, the distributions $\text{OE}_n(g)$, $\text{UE}_n(g)$ and $\text{SE}_n(g)$ for g one of the forms

$$e^{-x^2}, \quad x^\alpha e^{-x}, \quad x^a(1-x)^b. \quad (1-5)$$

These same weight functions occur in the theory of orthogonal polynomials [26]—they are associated with the three families of classical orthogonal polynomials Hermite, Laguerre and Jacobi respectively, and are themselves referred to as classical weight functions. The classical polynomials share many special properties not enjoyed by orthogonal polynomials associated with other weight functions. In the present study of matrix ensembles, we will see that the distributions $\text{OE}_n(g)$, $\text{UE}_n(g)$ and $\text{SE}_n(g)$ also have special features for g a classical weight function (1–5).

Our interest is in the properties of alternate eigenvalues in matrix ensembles. In particular we seek to determine the weights g for which alternate eigenvalues taken from a random union of two orthogonal ensembles form a unitary ensemble. Similarly we seek the weights g for which alternate eigenvalues from an orthogonal ensemble form a symplectic ensemble. The motivation for this study comes from recent work of Baik and Rains [4]. Consider the distribution $\text{OE}_n(e^{-x})$, n even, and order the eigenvalues $x_0 < x_1 < \cdots < x_{n-1}$. In [4] it was proved that after integrating out every second eigenvalue x_{n-1}, x_{n-3} , etc., the remaining eigenvalues have the distribution $\text{SE}_{n/2}(e^{-x})$. The proof of Baik and Rains is particular to the $a = 0$ case of the Laguerre ensemble. However other considerations lead these authors [5] to conjecture that in an appropriate scaled limit the distribution of the largest eigenvalue in the GSE corresponds to that of the second largest eigenvalue in the GOE. From this it is remarked that presumably the joint distribution of every second eigenvalue in the GOE coincides with the joint distribution of all the eigenvalues in the GSE, with an appropriate number of eigenvalues.

Baik and Rains [5] were also led to consider two GOE_n spectra, superimposing them at random, and integrating out every second eigenvalue of the resulting sequence. Results were presented which suggest that in the scaled $n \rightarrow \infty$ limit at the soft edge the distribution becomes that of GUE_∞ , appropriately scaled.

Such interrelationships between ensembles first occurred in the work of Dyson [8] on the circular ensembles of random unitary matrices. These ensembles have eigenvalue PDF proportional to

$$\prod_{0 \leq j < k < n} |e^{i\theta_k} - e^{i\theta_j}|^\beta, \quad 0 \leq \theta_j < 2\pi, \tag{1-6}$$

for $\beta = 1, 2,$ and 4 ($\text{COE}_n, \text{CUE}_n$ and CSE_n) respectively. Dyson conjectured that

$$\text{alt}(\text{COE}_n \cup \text{COE}_n) = \text{CUE}_n \tag{1-7}$$

which means that if two spectra from the COE_n distribution are superimposed at random with every second eigenvalue integrated out, the CUE_n distribution results. This was subsequently proved by Gunson [15]. Also, Mehta and Dyson [20] proved that integrating out every second eigenvalue from the distribution COE_n with n even gives the distribution $\text{CSE}_{n/2}$, or symbolically

$$\text{alt}(\text{COE}_n) = \text{CSE}_{n/2}. \tag{1-8}$$

The circular ensembles can be analyzed in the course of the present study of ensembles with real valued eigenvalues by making the stereographic projection

$$e^{i\theta_j} = \frac{1 - ix_j}{1 + ix_j}.$$

The PDF (1-6) then maps to

$$\prod_{j=0}^{n-1} \frac{1}{(1 + x_j^2)^{\beta(n-1)/2+1}} \prod_{0 \leq j < k < n} |x_k - x_j|^\beta$$

which is of the general type under consideration. Here the weight function is of the form

$$\frac{1}{(1 + x^2)^\alpha}, \quad \alpha > 1. \tag{1-9}$$

This only has a finite number of well defined moments and thus in this respect differs from the classical weight functions (1-5). On the other hand the corresponding orthogonal polynomials are $\{P_n^{(-\alpha, -\alpha)}(ix)\}_{n < \alpha - 1/2}$ [25], with $P_n^{(\alpha, \beta)}$ denoting the Jacobi polynomial, thus implying (1-9) can be viewed as a fourth classical weight function.

2. Pseudo-Ensembles

We begin with the orthogonal ensemble eigenvalue PDF (1–2), taking away the modulus sign, replacing n by l (to avoid overuse of the former) and rewriting the product as a determinant using the Vandermonde formula to obtain

$$\Delta(x) \prod_i g(x_i) = \det(x_i^j)_{0 \leq i, j < l} \prod_i g(x_i) = \det(g(x_i)x_i^j)_{0 \leq i, j < l}.$$

In particular, we note that each row corresponds to a variable, while each column corresponds to a function. Given a collection of n functions $F_i : \mathbb{R} \rightarrow \mathbb{R}$, we thus define the associated “orthogonal pseudo-ensemble” by the following “density”:

$$\det(F_j(x_i))_{0 \leq i, j < l}.$$

Thus any orthogonal ensemble is also an orthogonal pseudo-ensemble, but certainly not vice versa. Indeed, one has:

THEOREM 2.1. *Fix an integer $l > 0$, and let $G : \mathbb{R} \rightarrow \mathbb{R}$ be a function supported on at least n points. Then for a collection of n functions F_0, F_1, \dots, F_{l-1} , we have*

$$\det(F_j(x_i))_{0 \leq i, j < l} \propto \prod_i G(x_i) \Delta(x) \tag{2-1}$$

if and only if there exist l linearly independent polynomials p_i of degree at most $l - 1$ such that $F_i(x) = p_i(x)G(x)$ for all i and x .

PROOF. The “if” portion is easy enough:

$$\det(G(x_i)p_j(x_i))_{0 \leq i, j < l} = \prod_i G(x_i) \det(p_j(x_i))_{0 \leq i, j < l} \propto \prod_i G(x_i) \Delta(x),$$

since the polynomials are assumed linearly independent.

Now, suppose (2–1) holds. It will turn out to be convenient to restate the equation in terms of exterior products. Define a vector-valued function $V_F(x)$ by

$$V_F(x)_i = F_i(x).$$

Then we can write

$$\det(F_j(x_i))_{0 \leq i, j < l} = \left\langle V_F(x_0), \left(\bigwedge_{1 \leq i < l} V_F(x_i) \right) \right\rangle,$$

where $\langle \cdot, \cdot \rangle$ stands for the standard duality between 1-forms and $l - 1$ -forms. Consider this as a function of x_0 as the other variables range over the support of G ; we have:

$$V_F(x) \cdot \left(\bigwedge_{1 \leq i < l} V_F(x_i) \right) \propto G(x) \prod_{1 \leq i < l} (x - x_i).$$

Now, since G has at least l elements in its support, these functions span an l -dimensional space (this follows, for instance, from Lagrange’s interpolation

formula). On the other hand, the functions must clearly be linear combinations of the F_i . Since there only l functions F_i , it follows that we can write the F_i as linear combinations of the functions $G(x)x^i$, $0 \leq i < l$. But this is precisely what we wanted to prove. \square

Similarly, the density function of a symplectic ensemble can also be written as a determinant, namely

$$\Delta(x)^4 \prod_i g(x_i)^2 = \det(g(x_i)x_i^j, jg(x_i)x_i^{j-1})_{0 \leq i < l, 0 \leq j < 2l};$$

this follows by differentiating the Vandermonde determinant. When $\log(g)$ is differentiable, we can perform column transformations to put this determinant in the form

$$\det(F_j(x_i), F'_j(x_i))_{0 \leq i < l, 0 \leq j < 2l}; \tag{2-2}$$

simply take $F_j(x) = g(x)x^j$, and observe that

$$F'_j(x) - \frac{g'(x)}{g(x)}F_j(x) = jg(x)x^{j-1}.$$

In fact, we can often define (2-2) even when the functions F_j are not differentiable, by expressing it in terms of the 2-form-valued function

$$V_F^{(2)}(x) = \lim_{y \rightarrow x} \frac{1}{(x-y)} (V_F(x) \wedge V_F(y)).$$

For $F_j(x) = g(x)x^j$, we find that this is defined wherever g is continuous.

THEOREM 2.2. *Fix an integer $l > 1$, let O be a nonempty open subset of \mathbb{R} , and let $G : O \rightarrow \mathbb{R}$ be a continuous function supported on O . Then for a collection of continuous functions $F_j : O \rightarrow \mathbb{R}$ such that (2-2) is well-defined,*

$$\det(F_j(x_i), F'_j(x_i))_{0 \leq i < l, 0 \leq j < 2l} = \Delta(x)^4 \prod_i G(x_i)^2$$

on O^{2l} if and only if there exist linearly independent polynomials p_j of degree at most $2l - 1$ with $F_j(x) = G(x)p_j(x)$.

PROOF. Again the “if” case is straightforward. In the other direction, we can clearly divide each F_j by G , and thus may assume without loss of generality that $G = 1$ on O .

We first consider the case $l = 2$, for which

$$\det(F_j(x) F'_j(x) F_j(y) F'_j(y))_{0 \leq j < 4} = \langle V_F^{(2)}(x), V_F^{(2)}(y) \rangle = (x - y)^4.$$

As y varies over O , this spans a 5-dimensional function space; it follows that as y varies, $V_F^{(2)}(y)$ spans a 5-dimensional space (the dimension must be either 5 or 6; 6 clearly leads to a contradiction). In other words, there must be a linear dependence between the coefficients of $V_F^{(2)}(y)$. By replacing the F_i with an

orthogonal linear combination, we find that this dependence is without loss of generality of the form

$$V_F^{(2)}(y)_{01} = CV_F^{(2)}(y)_{23},$$

for some constant C . Now, if C were 0, then we would have

$$F_0(x)F_1'(x) = F_1(x)F_0'(x).$$

Now, let $I \subset O$ be an open interval in O . If either F_0 or F_1 were identically 0 on I , our determinant would be identically 0 on I^2 (contradiction); it follows that we may choose I so that both F_0 and F_1 are nonzero. Then we can divide both sides of the preceding equation by $F_0(x)F_1(x)$ and integrate; we find that $F_0 \propto F_1$ on I . But this again makes the determinant 0. We conclude that the linear dependence satisfied by $V_F^{(2)}$ must take the form

$$V_F^{(2)}(y)_{01} = CV_F^{(2)}(y)_{23}$$

with $C \neq 0$.

In particular, we find that the 2-form V' orthogonal to $V_F^{(2)}(y)$ is not itself in the span of $V_F^{(2)}(y)$. In particular, any 2-form can be written as a linear combination of V' and some of the $V_F^{(2)}(y)$. Taking the inner product with $V_F^{(2)}(x)$, we conclude that for $0 \leq i < j \leq 3$, we have

$$V_F^{(2)}(x)_{ij} = p_{ij}(x)$$

for some polynomial p of degree at most 4. Now, since $V_F(x) \wedge V_F^{(2)}(x) = 0$, we find:

$$F_0(x)p_{12}(x) - F_1(x)p_{02}(x) + F_2(x)p_{01}(x) = 0$$

for all i, j, k . Similarly, since

$$\frac{d}{dx}V_F^{(2)}(x) = V_F(x) \wedge V_F''(x),$$

we have

$$F_0(x)p'_{12}(x) - F_1(x)p'_{02}(x) + F_2(x)p'_{01}(x) = 0$$

Now, since $V_F^{(2)}(x)_{ki}$ is linearly independent of $V_F^{(2)}(x)_{ij}$, we can solve these two equations for F_1 and F_2 as rational multiples of F_0 , substitute into the equation $V_F^{(2)}(x)_{12} = p_{12}(x)$, then solve for F_0 . We find

$$F_0 = \frac{p_{01}p'_{02} - p_{02}p'_{01}}{\sqrt{D}}, \quad F_1 = \frac{p_{01}p'_{12} - p_{12}p'_{01}}{\sqrt{D}}, \quad F_2 = \frac{p_{02}p'_{12} - p_{12}p'_{02}}{\sqrt{D}},$$

where

$$D = \det \begin{pmatrix} p_{01} & p_{12} & p_{20} \\ p'_{01} & p'_{12} & p'_{20} \\ p''_{01} & p''_{12} & p''_{20} \end{pmatrix}.$$

We observe that each numerator has degree at most 6, as does the polynomial D . In particular, if we exclude any given F , we can express the squares of the other F as rational functions with common denominator of degree at most 6. It

follows that the functions F^2 have at most 8 poles between them, and thus that we can write

$$\begin{aligned} F_0(x) &= p_0(x)p(x)^{-1/2}, & F_1(x) &= p_1(x)p(x)^{-1/2}, \\ F_2(x) &= p_2(x)p(x)^{-1/2}, & F_3(x) &= p_3(x)p(x)^{-1/2}, \end{aligned}$$

where $p_0, p_1, p_2,$ and p_3 are polynomials of degree at most 7 and p is a polynomial of degree at most 8.

We now need to show that, in fact, each F_i is a polynomial of degree at most 3. By the usual factorization, we find:

$$\det(p_j(x) p'_j(x) p_j(y) p'_j(y))_{0 \leq j < 4} \propto p(x)p(y)(x - y)^4, \tag{2-3}$$

valid on \mathbb{R} . Without loss of generality, we may assume that the constant of proportionality is 1, and that $p(0) = 1$. Dividing both sides by $(x - y)^4$ and taking the limit as $x, y \rightarrow 0$, we find:

$$\det \begin{pmatrix} p_0(0) & p_1(0) & p_2(0) & p_3(0) \\ p'_0(0) & p'_1(0) & p'_2(0) & p'_3(0) \\ p''_0(0) & p''_1(0) & p''_2(0) & p''_3(0) \\ p'''_0(0) & p'''_1(0) & p'''_2(0) & p'''_3(0) \end{pmatrix} = 1$$

Applying a suitable linear transformation to the polynomials p_i , we have, without loss of generality,

$$\begin{aligned} p_0(x) &= 1 + p_{04}x^4 + p_{05}x^5 + p_{06}x^6 + p_{07}x^7, \\ p_1(x) &= x + p_{14}x^4 + p_{15}x^5 + p_{16}x^6 + p_{17}x^7, \\ p_2(x) &= x^2 + p_{24}x^4 + p_{25}x^5 + p_{26}x^6 + p_{27}x^7, \\ p_3(x) &= x^3 + p_{34}x^4 + p_{35}x^5 + p_{36}x^6 + p_{37}x^7. \end{aligned}$$

We can then solve for $p(x)$ by taking $y = 0$ above; we find

$$p(x) = x^{-4}(p_2(x)p'_3(x) - p_3(x)p'_2(x)).$$

At this point, we can compare coefficients on both sides of (2-3), obtaining a number of polynomial equations relating the coefficients p_{ij} , where $0 \leq i \leq 3$ and $4 \leq j \leq 7$. The resulting ideal can be verified (using Magma, for instance) to contain the polynomials $(p_{25} + p_{34}p_{35} - p_{36})^2$, $(p_{26} + p_{34}p_{36} - p_{37})^2$, and $(p_{27} + p_{34}p_{37})^2$; passing to the radical, we can then solve for p_{ij} , $0 \leq i \leq 2$, $4 \leq j \leq 7$. Substituting in, we find that $p(x)$ is now a square, and that each $p_i(x)$ is a multiple of $\sqrt{p(x)}$. In other words, each $F_i(x)$ is a polynomial of degree at most 3, and we are done with the case $l = 2$.

It remains only to show that we can reduce the cases $l > 2$ to cases of lower dimension. Choose a particular element $x_0 \in O$. By replacing the F_i with appropriate linear transformations, we may assume

$$V_F^{(2)}(x_0) = Ce_1 \wedge e_2,$$

for some nonzero constant C . In particular, we find that

$$\begin{aligned} \det(F_j(x_i))_{0 \leq i < l, 0 \leq j \leq 2l} \\ = C \det(F_j(x_i))_{1 \leq i < l, 2 \leq j \leq 2l} \propto G(x_0) \Delta(x_1, x_2, \dots, x_{l-1})^4 \prod_{1 \leq i < l} x_i^4 G(x_i)^2. \end{aligned}$$

By induction, it follows that for $2 \leq i < l$, there exist polynomials $p_i(x)$ of degree at most $2l - 1$ such that $F_i(x) = (x - x_0)^2 G(x) p_i(x)$ on O . Undoing our linear transformations, we find that for every polynomial $p(x)$ of degree at most $2l - 1$ vanishing to second order at x_0 , we can write $G(x)p(x)$ as a linear combination of the $F_i(x)$. But this was independent of our choice of x_0 . In particular, taking x'_0 to be any other element of O , we have

$$1 = \frac{3(x_0 - x'_0)(x - x_0)^2 + 2(x - x_0)^3 + 3(x_0 - x'_1)(x - x'_0)^2 - 2(x - x'_0)^3}{(x_0 - x'_0)^3}$$

and

$$(x - x_0) = \frac{-2(x_0 - x'_0)(x - x_0)^2 - (x - x_0)^3 - (x_0 - x'_0)(x - x'_0)^2 + (x - x'_0)^3}{(x_0 - x'_0)^2}.$$

It follows that for any polynomial $p(x)$ of degree at most $2l - 1$, $G(x)p(x)$ is a linear combination of the $F_i(x)$. By dimensionality, it follows that each $F_i(x)$ is itself of the form $G(x)p(x)$, and we are done. \square

3. Linear Fractional Transformations

It will be convenient in the sequel to determine how matrix ensembles behave under a linear fractional transformation (LFT) or change of variables. To be precise, let f be a weight function, and consider what the density of one of its associated matrix ensembles is in terms of the variables y_i defined by $x_i = (\alpha y_i + \beta)/(\gamma y_i + \delta)$. Clearly, we need only determine how $\Delta(x)$ and $\prod_i dx_i$ transform.

We readily compute:

$$dx = \frac{\alpha\delta - \beta\gamma}{(\gamma y + \delta)^2},$$

thus answering that question. As for Δ :

LEMMA 3.1. *Let y_0, y_1, \dots, y_{l-1} be a collection of l real numbers. Then for any α, β, γ and δ such that $\gamma y_i + \delta$ is never 0,*

$$\Delta \left(\frac{\alpha y_i + \beta}{\gamma y_i + \delta} \right) = (\alpha\delta - \beta\gamma)^{l(l-1)/2} \prod_i (\gamma y_i + \delta)^{1-l} \Delta(y_i).$$

PROOF. For each $i < j$, we have

$$\frac{\alpha y_j + \beta}{\gamma y_j + \delta} - \frac{\alpha y_i + \beta}{\gamma y_i + \delta} = \frac{\alpha\delta - \beta\gamma}{(\gamma y_i + \delta)(\gamma y_j + \delta)} (y_j - y_i).$$

Multiplying over $i < j$, we are done. \square

We thus obtain the following transformation rules:

THEOREM 3.2. *Let f be any weight function. Under the change of variables $x_i = (\alpha y_i + \beta)/(\gamma y_i + \delta)$, we have*

$$\begin{aligned} \text{OE}_l(f(x)) &\rightarrow \text{OE}_l(|\alpha\delta - \beta\gamma|^{(l+1)/2}(\gamma y + \delta)^{-1-l} \tilde{f}(y)), \\ \text{UE}_l(f(x)) &\rightarrow \text{UE}_l(|\alpha\delta - \beta\gamma|^l(\gamma y + \delta)^{-2l} \tilde{f}(y)), \\ \text{SE}_l(f(x)) &\rightarrow \text{SE}_l(|\alpha\delta - \beta\gamma|^{2l-1}(\gamma y + \delta)^{2-4l} \tilde{f}(y)), \end{aligned}$$

where $\tilde{f}(y) = f((\alpha y + \beta)/(\gamma y + \delta))$, the normalization constants are the same on both sides, and for OE_{2l} , $\gamma y + \delta$ must be positive over the support of \tilde{f} .

PROOF. When $\alpha\delta - \beta\gamma < 0$, the LFT reverses the order of integration, thus justifying the extra factor of $(-1)^l$ introduced for UE_l and SE_l . For OE_l , there is a more subtle difficulty, namely that the relative order of the eigenvalues is significant, and can change. If we simply reverse the order, this is not a problem (the total effect is $(-1)^{l(l+1)/2}$, thus cancelling out the sign of $\alpha\delta - \beta\gamma$). So we can restrict to the case $\alpha\delta - \beta\gamma > 0$. The effect of the LFT is then to cyclically shift the ordering, taking the eigenvalues with $x > \alpha/\gamma$ and making them smallest. If there are k such eigenvalues, the sign of the Vandermonde matrix is changed by $(-1)^{k(l-1)}$; thus if l is odd, there is no problem. On the other hand, if l is even, we have a problem unless the eigenvalues are restricted to only one side of α/γ , or equivalently that $\gamma y + \delta$ has constant sign over the support of \tilde{f} . Since

$$\frac{\alpha y + \beta}{\gamma y + \delta} = \frac{-\alpha y - \beta}{-\gamma y - \delta},$$

we may take this sign to be positive. □

REMARK. For algebraic purposes, we can often ignore the constraint $\gamma y + \delta > 0$, since the transform still has the correct form to be a matrix ensemble density, despite not being nonnegative.

The upshot of this is that we can use this freedom to send a suitably chosen point to ∞ , thus simplifying our analysis below.

4. The Main Results

For a matrix ensemble M , we define $\text{even}(M)$ to be the ensemble obtained by taking the second largest, fourth largest, etc. eigenvalues of M , and similarly for $\text{odd}(M)$.

When considering $\text{even}(M)$ or $\text{odd}(M)$ in the cases $M = \text{OE}_n \cup \text{OE}_n$ and $M = \text{OE}_n \cup \text{OE}_{n+1}$, the following lemma is crucial:

LEMMA 4.1. *For any integer $n > 0$,*

$$\sum_{\substack{S \subset \{0,1,\dots,2n-1\} \\ |S|=n}} \Delta(x_S) \Delta(x_{\{0,1,\dots,2n-1\}-S}) = 2^n \Delta(x_{\{0,2,\dots,2n-2\}}) \Delta(x_{\{1,3,\dots,2n-1\}})$$

and

$$\sum_{\substack{S \subset \{0,1,\dots,2n\} \\ |S|=n+1}} \Delta(x_S) \Delta(x_{\{0,1,\dots,2n\}-S}) = 2^n \Delta(x_{\{0,2,\dots,2n\}}) \Delta(x_{\{1,3,\dots,2n-1\}})$$

PROOF. Consider what happens when we exchange x_i and x_{i+2} in a term of either equation. If $i, i + 2 \in S$ or $i, i + 2 \notin S$, then

$$\Delta(x_S) \Delta(x_{\{0,1,\dots,l-1\}-S}) \rightarrow -\Delta(x_S) \Delta(x_{\{0,1,\dots,l-1\}-S}),$$

since Δ is alternating. Otherwise, we see that every factor $x_j - x_k$ with $j > k$ is taken to another such factor, *except* for the factor $x_{i+1} - x_i$ or $x_{i+2} - x_{i+1}$, whichever is present. So each term in our sum is taken to the negative of a term from our sum; it follows that the sum is alternating under parity-preserving permutations. It follows that it must be a multiple of

$$\Delta(x_{\{0,2,\dots,2\lfloor l/2\rfloor-2\}}) \Delta(x_{\{1,3,\dots,2\lfloor l/2\rfloor-1\}}).$$

By degree considerations, it remains only to verify the constant, which we can do by considering the coefficient of largest degree in x_0 , and applying induction. \square

REMARK. The even case of this lemma is implicit in [15], where it was used to analyze $\text{even}(\text{OE}_n \cup \text{OE}_n)$ with respect to the weight function 1 on the unit circle.

From the lemma, it follows that the density of $\text{OE}_n(f) \cup \text{OE}_n(f)$, expressed in terms of ordered variables, is proportional to

$$\prod_{0 \leq i \leq 2n} f(x_i) \Delta(x_{\{0,2,\dots,2n-2\}}) \Delta(x_{\{1,3,\dots,2n-1\}});$$

similarly, the density of $\text{OE}_n(f) \cup \text{OE}_{n+1}(f)$ is proportional to

$$\prod_{0 \leq i \leq 2n} f(x_i) \Delta(x_{\{0,2,\dots,2n\}}) \Delta(x_{\{1,3,\dots,2n-1\}}).$$

For some weight functions f , if we integrate over the odd/even variables, the resulting density is the density of a unitary ensemble; we wish to determine precisely when that is. We first consider the case $\text{even}(\text{OE}_2(f) \cup \text{OE}_3(f))$.

THEOREM 4.2. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function which is differentiable on a possibly unbounded open interval $I \subset \mathbb{R}$ and 0 elsewhere. Suppose $\text{even}(\text{OE}_2(f) \cup \text{OE}_3(f)) = \text{UE}_2(g)$ for some function g . Then up to a linear transformation of variables, f must have one of the following forms. On the interval $(0, 1)$:*

$$\begin{aligned} f(x) &\propto x^\alpha (1-x)^\beta (1-rx)^{-4-\alpha-\beta}, & \alpha, \beta > -1, r < 1 \\ f(x) &\propto x^{-4-\alpha} e^{-1/x} (1-x)^\alpha, & \alpha > -1. \end{aligned}$$

On the interval $(0, \infty)$:

$$\begin{aligned} f(x) &\propto x^\alpha(1-rx)^\beta, & \alpha > -1, \alpha + \beta < -3, r < 0, \\ f(x) &\propto x^{-4-\alpha}e^{-1/x}, & \alpha > -1, \\ f(x) &\propto x^\alpha e^{-x}, & \alpha > -1. \end{aligned}$$

Finally, on the entire real line:

$$\begin{aligned} f(x) &\propto (1+x^2)^\alpha, & \alpha < -\frac{3}{2} \\ f(x) &\propto e^{-x^2/2}. \end{aligned}$$

PROOF. We need to integrate this over the variables x_{2i} , and thus need to evaluate the determinant

$$\det \left(\int_{[x_{2i-1}, x_{2i+1}]} f(x)x^j dx \right)_{0 \leq i, j \leq 2},$$

where we take $x_{-1} = a$ to be the left endpoint of I , and $x_5 = b$ to be the right endpoint of I . In particular, we need to determine when there exists a function $g(x)$ with

$$\det \left(\int_{[x_{2i-1}, x_{2i+1}]} f(x)x^j dx \right)_{0 \leq i, j \leq 2} \propto g(x_1)g(x_3)(x_3 - x_1).$$

As in [19, Section 10.6], we may use row operations to transform this to

$$\det (F_j(x_{2i+1}))_{0 \leq i, j \leq 2},$$

where we define

$$F_j(y) = \int_{[a, y]} f(x)x^j dx.$$

We cannot quite apply Theorem 2.1, however, since the last column of our determinant is constant. However, we clearly have $F_0(b) > 0$, so we can eliminate that column, obtaining

$$F_0(b) \det \left(F_j(x_{2i+1}) - \frac{F_j(b)}{F_0(b)} F_0(x_{2i+1}) \right)_{0 \leq i, j \leq 1}.$$

This, then, satisfies the hypotheses of Theorem 2.1; there thus exist linear polynomials p_j such that

$$p_2(x)(F_1(x) - C_1 F_0(x)) = p_1(x)(F_2(x) - C_2 F_0(x)), \tag{4-1}$$

where we have set

$$C_i = \frac{F_i(b)}{F_0(b)}.$$

Differentiating twice and using the definition of F_i , we find

$$\begin{aligned} &(p_2(x)(x - C_1) - p_1(x)(x^2 - C_2)) f'(x) \\ &= (-2(x - C_1)p_2'(x) + 2(x^2 - C_2)p_1'(x) - p_2(x) + 2xp_1(x)) f(x). \end{aligned}$$

We can thus solve this for $f'(x)/f(x)$; we find that $f'(x)/f(x)$ has the form $p(x)/q(x)$ with $\deg(p) \leq 2$, $\deg(q) \leq 3$, and $\deg(xp + 4q) \leq 2$. We observe that these conditions are, naturally, preserved by linear fractional transformations. In particular, by applying a suitable linear fractional transformation, we may insist that q be strictly cubic, and that both endpoints of I be finite (possibly equal). (The result may very well no longer be a matrix ensemble, but as we noted above, this does not affect any algebraic conclusions.)

Now, consider how $f(x)$ and $q(x)$ must behave at 0 and 1. Differentiating (4-1) once and taking a limit $x \rightarrow x_{-1}$ we find, since each $F_i(x_{-1}) = 0$,

$$\lim_{x \rightarrow x_{-1}} (p_2(x)(x - C_1) + p_1(x)(C_2 - x))f(x) = 0$$

But this is just $\lim_{x \rightarrow x_{-1}} q(x)f(x)$. If $q(x_{-1}) \neq 0$, we must have $\lim_{x \rightarrow x_{-1}} f(x) = 0$. Then

$$\lim_{x \rightarrow x_{-1}} \frac{f'(x)}{f(x)} = \infty.$$

The only way this can happen is if $q(x_{-1}) = 0$ after all. Similarly, we have $q(x_{2n+1}) = 0$.

Suppose first that $a \neq b$. Then up to LFT, we may insist that $a = 0$ and $b = 1$, and thus $q(0) = q(1) = 0$. We thus have two possibilities. The first is that $q(x)$ has an additional zero, neither 0 nor 1. In this case, integrating f'/f and taking into account the constraints on $p(x)$, we obtain

$$f(x) = x^\alpha(1-x)^\beta(1-rx)^{-4-\alpha-\beta}$$

Now, for $\int f(x)$ not to diverge at 0, we must have $\alpha > -1$, and similarly $\beta > -1$. But then $-4-\alpha-\beta < -2$; it follows that $(1-rx)$ must be nonzero on $(0, 1)$; in particular, $r < 1$. The other possibility is that $q(x)$ has a double root, without loss of generality, at 0. Upon integrating f'/f , we obtain

$$f(x) = x^{-4-\alpha} e^{-\beta/x} (1-x)^\alpha,$$

and find $\alpha > -1$, $\beta > 0$. The possibilities for $(0, \infty)$ then follow by LFT.

The other case we must consider is $I = \mathbb{R}$, and thus $\deg(q) = 2$, $\deg(p) = 1$. If q had a simple root in \mathbb{R} , it would have two (say 0 and 1, without loss of generality), and thus f would have the form $x^\alpha(1-x)^\beta$ with $\alpha, \beta > -1$. But then the integral for F_2 would diverge. Similarly, if q had a double root at 0, f would have the form $x^\alpha e^{-\beta/x}$, which would diverge on one side of 0. Thus either q has a pair of complex roots, or $\deg(q) = 0$. In the first case, a linear transformation takes the roots to $\pm i$, and thus

$$f(x) = (1+x^2)^\alpha.$$

For F_2 to be well-defined, we must have $\alpha < -3/2$. The other possibility gives $\log(f(x)) = ax^2 + bx + c$; thus a linear transformation gives

$$f(x) = e^{-x^2/2}. \quad \square$$

We can now extend this to $n \geq 2$.

THEOREM 4.3. *Fix an integer $n \geq 2$. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function which is differentiable on a possibly unbounded open interval $I \subset \mathbb{R}$ and 0 elsewhere. Suppose $\text{even}(\text{OE}_n(f) \cup \text{OE}_{n+1}(f)) = \text{UE}_n(g)$ for some function g . Then up to a linear transformation of variables, f and g can have precisely the following forms. On the interval $(0, 1)$:*

$$\begin{aligned} f(x) &\propto x^\alpha(1-x)^\beta(1-rx)^{-n-2-\alpha-\beta}, \quad \alpha, \beta > -1, r < 1, \\ g(x) &\propto x^{2\alpha+1}(1-x)^{2\beta+1}(1-rx)^{-2n-3-2\alpha-2\beta}, \\ f(x) &\propto x^{-n-2-\alpha}e^{-1/2x}(1-x)^\alpha, \quad \alpha > -1, \\ g(x) &\propto x^{-2n-2-2\alpha}e^{-1/x}(1-x)^{2\alpha+1}. \end{aligned}$$

On the interval $(0, \infty)$:

$$\begin{aligned} f(x) &\propto x^\alpha(1-rx)^\beta, \quad \alpha > -1, \alpha + \beta < -n - 1, r < 0, \\ g(x) &\propto x^{2\alpha+1}(1-rx)^{2\beta+1}, \\ f(x) &\propto x^{-n-2-\alpha}e^{-1/2x}, \quad \alpha > -1, \\ g(x) &\propto x^{-2n-2-2\alpha}e^{-1/x}, \\ f(x) &\propto x^\alpha e^{-x/2}, \quad \alpha > -1, \\ g(x) &\propto x^{2\alpha+1}e^{-x}. \end{aligned}$$

Finally, on the entire real line:

$$\begin{aligned} f(x) &\propto (1+x^2)^\alpha, \quad \alpha < -(n+1)/2, \\ g(x) &\propto (1+x^2)^{2\alpha+1}, \\ f(x) &\propto e^{-x^2/2}, \\ g(x) &\propto e^{-x^2}. \end{aligned}$$

PROOF. As for $n = 2$, the issue is when

$$\det(F_j(x_{2i+1}))_{0 \leq i, j \leq n}$$

takes the form of an orthogonal ensemble. Applying an LFT as necessary, we may assume that $a = -\infty$. Now differentiate with respect to x_1 , divide by $x_1^{n-1}f(x_1)$, and take a limit as $x_1 \rightarrow -\infty$. On the one hand, this operation takes orthogonal ensembles to orthogonal ensembles. On the other hand, we can then expand along the first column, finding that

$$\det(F_{j-1}(x_{2i+1}))_{1 \leq i, j \leq n}$$

must take the form of an orthogonal ensemble. By induction, we find that $f(x)$ must satisfy the constraints valid for $n - 1$. Upon undoing the LFT, we obtain the desired “only if” result.

It remains to show that each of the above weight functions actually do work. We need only consider the following possibilities:

$$\begin{aligned} f(x) &= x^\alpha(1-x)^\beta && \text{on } (0, 1), \\ f(x) &= x^\alpha e^{-x} && \text{on } (0, \infty), \\ f(x) &= (1+x^2)^\alpha && \text{on } \mathbb{R}, \\ f(x) &= e^{-x^2/2} && \text{on } \mathbb{R}, \end{aligned}$$

since the others are all images of these under LFTs.

For $f(x) = x^\alpha(1-x)^\beta$, observe that

$$(\alpha + \beta + j + 2)F_{j+1}(x) - (\alpha + j + 1)F_j(x) = -x^j(x^{\alpha+1}(1-x)^{\beta+1});$$

this is true for $x = 0$, and both sides have the same derivative. In particular, for each j , we have a polynomial $p_j(x)$ of degree $j - 1$ and a constant C_j with

$$F_j(x) = C_j F_0(x) + p_j(x)(x^{\alpha+1}(1-x)^{\beta+1}).$$

In particular, this must be true for $x = 1$, and thus $C_j = F_j(1)/F_0(1)$ as required. We thus find that we obtain a unitary ensemble with weight function proportional to $x^{2\alpha+1}(1-x)^{2\beta+1}$. Similarly, for $f(x) = x^\alpha e^{-x}$, we have

$$F_{j+1}(x) - (\alpha + j + 1)F_j(x) = -x^{\alpha+j+1}e^{-x},$$

so $g(x) \propto x^{2\alpha+1}e^{-2x}$.

For $f(x) = (1+x^2)^\alpha$, we find

$$(2\alpha + j + 2)F_{j+1}(x) + jF_{j-1} = x^j(1+x^2)^{\alpha+1}$$

This allows us to solve for each F_j except F_0 ; we obtain $g(x) \propto (1+x^2)^{2\alpha+1}$. Finally, for $f(x) = e^{-x^2/2}$, we have:

$$F_{j+1} - jF_{j-1} = -x^j e^{-x^2/2},$$

and $g(x) \propto e^{-x^2}$. □

REMARK. We observe that in each case $g(x) \propto f(x)^2 q(x)$.

For $\text{even}(\text{OE}_n \cup \text{OE}_n)$, the calculations are analogous, and we have:

THEOREM 4.4. *Fix an integer $n \geq 2$. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function which is differentiable on a possibly unbounded open interval $I \subset \mathbb{R}$ and 0 elsewhere. Suppose $\text{even}(\text{OE}_n(f) \cup \text{OE}_n(f)) = \text{UE}_n(g)$ for some function g . Then up to an order-preserving linear transformation of variables, f and g must have one of the following forms. On an interval with right endpoint 0:*

$$\begin{aligned} f(x) &\propto (-x)^\alpha(1-rx)^{-n-1-\alpha}, && \alpha > -1, 1 \notin rI, \\ g(x) &\propto (-x)^{2\alpha+1}(1-rx)^{-2n-1-2\alpha}, \\ f(x) &\propto (-x)^{-n-1}e^{1/x}, \\ g(x) &\propto (-x)^{-2n}e^{2/x}. \end{aligned}$$

On an interval of the form (a, ∞) , $a > -\infty$:

$$\begin{aligned} f(x) &\propto (1 - rx)^\alpha, & \alpha < -n, r < 0, \\ g(x) &\propto (1 - rx)^{2\alpha+1}, \\ f(x) &\propto e^{-x}, \\ g(x) &\propto e^{-2x} \end{aligned}$$

On the entire real line, no possibilities exist.

REMARK. The relation between g and f is here slightly modified, by removing the factor of g corresponding to the left endpoint. For $\text{odd}(\text{OE}_n \cup \text{OE}_n)$, we remove the factor corresponding to the right endpoint, and for $\text{odd}(\text{OE}_{n-1} \cup \text{OE}_n)$, we remove both factors.

For $\text{odd}(\text{OE}_n \cup \text{OE}_n)$, we simply reverse the ordering. For $\text{odd}(\text{OE}_n \cup \text{OE}_{n+1})$, we have:

THEOREM 4.5. Fix an integer $n \geq 2$. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function which is differentiable on a possibly unbounded open interval $I \subset \mathbb{R}$ and 0 elsewhere. Suppose $\text{odd}(\text{OE}_{n-1}(f) \cup \text{OE}_n(f)) = \text{UE}_n(g)$ for some function g . Then f and g have the form

$$\begin{aligned} f(x) &\propto (1 - rx)^{-n}, \\ g(x) &\propto (1 - rx)^{-2n+1} \end{aligned}$$

for some r (possibly ∞) with $1/r \notin I$.

PROOF. The only tricky aspect of this case is that the determinant we must analyze is no longer of the form to which Theorem 2.1 applies; to be precise, we need

$$\det(F_j(x_{2i+2}) - F_j(x_{2i}))_{0 \leq i < n}$$

to have orthogonal ensemble form. But this determinant is clearly equal to the determinant of the block matrix

$$\begin{pmatrix} (1) & (0)_{0 \leq i < n} \\ (F_j(x_0))_{0 \leq j < n} & (F_j(x_{2i+2}) - F_j(x_{2i}))_{0 \leq i, j < n} \end{pmatrix}.$$

Adding the first column to the other columns, we can then apply Theorem 2.1, and argue as above. \square

We finally consider a fifth possibility for decimation. Recall that for the circular ensemble results cited above, while there was a *local* notion of order, there was no notion of largest. This suggests that we consider the ensemble derived by choosing randomly between $\text{odd}(M)$ and $\text{even}(M)$. More precisely, for an ensemble with an even number of variables, we define $\text{alt}(M)$ to be $\text{even}(M)$ with probability $\frac{1}{2}$ and $\text{odd}(M)$ with probability $\frac{1}{2}$.

THEOREM 4.6. *Fix an integer $n \geq 2$. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function which is differentiable on a possibly unbounded open interval $I \subset \mathbb{R}$ and 0 elsewhere. Suppose $\text{alt}(\text{OE}_n(f) \cup \text{OE}_n(f)) = \text{UE}_n(g)$ for some function g . Then up to a linear transformation of variables, f and g have the form*

$$\begin{aligned} f &= (1 + x^2)^{-(n+1)/2}, \\ g &= (1 + x^2)^{-n}. \end{aligned}$$

PROOF. Consider the determinants associated to $\text{even}(\text{OE}_n(f) \cup \text{OE}_n(f))$ and $\text{odd}(\text{OE}_n(f) \cup \text{OE}_n(f))$. Up to cyclic shift, only one column differs between the two determinants, thus allowing us to express their sum as a determinant. When n is even, the ‘special’ column takes the form

$$(F_j(x_0) + F_j(x_{2n-1}) - F_j(I))_{0 \leq j < n};$$

here $F_j(I) = \int_{x \in I} x^j f(x)$. Taking appropriate linear combinations, we obtain the determinant

$$\det(F_j(x_i) - F_j(I)/2)_{0 \leq i, j < n}.$$

When n is odd, the special column takes the form

$$(F_j(x_{2n-1}) - F_j(x_0) - F_j(I))_{0 \leq j < n};$$

this leads (up to sign) to the $n + 1 \times n + 1$ block determinant

$$\det \begin{pmatrix} 0 & (F_j(I))_{0 \leq j < n} \\ 1 & (F_j(x_i))_{0 \leq i, j < n} \end{pmatrix}.$$

We first analyze the case n odd. In this case, the usual theory tells us that there exist polynomials $p_j(x)$ and $q(x)$ of degree at most $n - 1$ with

$$F_j(x) - C_j F_0(x) = p_j(x)/q(x)$$

for all j , with C_j as above. Now, evaluating this at an endpoint of I , we find that the polynomials p_j must have a common root (possibly ∞). In particular, it follows that f must satisfy the conditions of Theorem 4.3. On the other hand, we find that

$$x^j f(x) - C_j f(x) = \frac{d}{dx} \frac{p_j(x)}{q(x)}$$

for each j ; in particular, $f(x)$ must be a rational function. We therefore have the following possibilities to consider:

$$\begin{aligned} f(x) &= x^\alpha (1 - x)^\beta && \text{for } \alpha, \beta \in \mathbb{N}, \\ f(x) &= (1 + x^2)^{-\alpha} && \text{for } \alpha \in \mathbb{N}, \alpha > n/2, \end{aligned}$$

on $(0, 1)$ and \mathbb{R} respectively. In the first case, we find that each

$$F_j(x) - C_j F_0(x)$$

is a polynomial of degree $j + 2$. In particular, $F_{n-1}(x) - C_{n-1} F_0(x)$ is a polynomial of degree $n + 1$, contradicting the bound on $\deg(p_j(x))$.

In the second case, we observe that

$$F_j(x) - C_j F_0(x) = r_j(x)(1 + x^2)^{1-\alpha}$$

for polynomials $r_j(x)$ of degree $j - 1$. In particular, we find that $p_1(x)/q(x) \propto (1 + x^2)^{1-\alpha}$, implying, since $\alpha > n/2 > 1$, that

$$q(x) \propto (1 + x^2)^{\alpha-1}$$

Since $\deg(q) \leq n - 1$, we have $n/2 < \alpha \leq (n + 1)/2$, the only integral solution of which is $\alpha = (n + 1)/2$. In this case, the relevant degree bounds all hold, and thus the determinant is indeed of the correct form, giving $g(x) \propto (1 + x^2)^{1-2\alpha} = (1 + x^2)^{-n}$ as required.

For n even, we must have polynomials p_i of degree at most $n - 1$ with

$$p_i(x)(F_j(x) - C'_j) = p_j(x)(F_i(x) - C'_i),$$

where we write C'_j for $F_j(I)/2$. We can rewrite this as:

$$p_0(x)(F_j(x) - C_j F_0(x)) = (p_j(x) - C_j p_0(x))(F_0(x) - C'_0),$$

using the fact that $C_j = C'_j/C'_0$. For $n > 2$, we conclude that $f(x)$ must satisfy the conditions of Theorem 4.3. Now, if the endpoints of I are different, then we find, since $F_0(x) - C'_0 = \pm C'_0$ at both endpoints, that each $p_j(x) - C_j p_0(x) = 0$ at both endpoints. But this causes the polynomials to be linearly dependent, a contradiction. On the other hand, in the other cases, we know that $C_1 = 0$ and $p_1 \propto 1$. In both cases, we obtain from the identity for $F_1(x) - C_1 F_0(x)$ a differential equation for $p_0(x)$. For $e^{-x^2/2}$, no polynomial solution to the equation exists. For $(1 + x^2)^{-\alpha}$, we can find an explicit power series solution to the equation, and find that a polynomial solution exists only when α is half-integral, when the solution has degree $2\alpha - 2$. As above, this leaves only one possibility for α , namely $\alpha = (n + 1)/2$, as required.

It remains to consider $n = 2$. Here we can twice differentiate the equation

$$p_0(x)(F_1(x) - C'_1) = p_1(x)(F_0(x) - C'_0)$$

(with p_0 and p_1 linear) to deduce that

$$f'(x)/f(x) = p(x)/q(x)$$

with $\deg(p) \leq 1$, $\deg(q) \leq 2$, and $\deg(xp + 3q) \leq 1$. So up to LFT, $f(x)$ must have one of the forms

$$\begin{aligned} f(x) &= x^\alpha && \text{for } \alpha \geq -\frac{3}{2}, \\ f(x) &= e^{-x}, \\ f(x) &= (1 + x^2)^{-3/2}. \end{aligned}$$

(note that if we exchange 0 and ∞ in the first case, we replace α with $-3 - \alpha$, justifying our restriction on α .) In the first case, if $\alpha \neq -1$, the following:

$$\frac{x^{\alpha+2} - D_1}{x^{\alpha+1} - D_0}$$

must be a linear polynomial for suitable constants D_1 and D_0 respectively proportional to C'_1 and C'_0 (and thus $D_0 \neq 0$). We deduce therefore that $\alpha = 0$. But then we readily determine that only the empty interval satisfies the requirements. For x^{-1} and e^{-x} , there are not even appropriate choices for D_0 and D_1 (since both $\log(x)$ and e^{-x} are transcendental functions). Finally, for the third choice, we readily verify that decimation indeed works as required. \square

We now turn our attention to decimations of single orthogonal ensembles. We have, quite simply:

THEOREM 4.7. *Fix an integer $n \geq 2$. For any functions f and g with f differentiable on a possibly unbounded open interval $I \subset \mathbb{R}$ and 0 elsewhere, the following equivalences hold (where we write $\text{OE}_{2n} f = \text{OE}_{2n}(f)$ and so on):*

$$\text{even}(\text{OE}_{2n} f \cup \text{OE}_{2n+1} f) = \text{UE}_{2n} g \iff \text{even}(\text{OE}_{2n+1} f) = \text{SE}_n((g/f)^2), \quad (4-2)$$

$$\text{even}(\text{OE}_{2n} f \cup \text{OE}_{2n} f) = \text{UE}_{2n} g \iff \text{even}(\text{OE}_{2n} f) = \text{SE}_n((g/f)^2), \quad (4-3)$$

$$\text{odd}(\text{OE}_{2n} f \cup \text{OE}_{2n} f) = \text{UE}_{2n} g \iff \text{odd}(\text{OE}_{2n} f) = \text{SE}_n((g/f)^2), \quad (4-4)$$

$$\text{odd}(\text{OE}_{2n-1} f \cup \text{OE}_{2n} f) = \text{UE}_{2n} g \iff \text{odd}(\text{OE}_{2n-1} f) = \text{SE}_n((g/f)^2), \quad (4-5)$$

$$\text{alt}(\text{OE}_{2n} f \cup \text{OE}_{2n} f) = \text{UE}_{2n} g \iff \text{alt}(\text{OE}_{2n} f) = \text{SE}_n((g/f)^2). \quad (4-6)$$

PROOF. Consider, for instance, $\text{even}(\text{OE}_{2n+1}(f))$. Once we integrate along the largest, third largest, etc., variables and do some simplification, the resulting matrix has columns $(F_j(x_{2i+1}))_{0 \leq j \leq 2n}$ and $(x_{2i+1}^j f(x_{2i+1}))_{0 \leq j \leq 2n}$, with the last column given by $(F_j(b))_{0 \leq j \leq 2n}$. In particular, we note that aside from the last, constant, column, the columns come in pairs, one the derivative of the other. Thus the determinant is essentially of the form considered in Theorem 2.2. In particular, it has a symplectic ensemble form if and only if the determinant

$$\det(F_j(x_{i+1}))_{0 \leq i, j \leq 2n}$$

(of which our determinant is a derivative) has an orthogonal ensemble form. But by the proof of Theorems 4.2 and 4.3, this precisely what we needed to show. (The statement about the resulting weight functions is straightforward.) Similar arguments apply for the remaining equivalences. \square

5. Random matrix applications

In random matrix applications f , g and $(g/f)^2$ must be (up to the scale of x) one of the four classical forms (1–5) and (1–9). So specializing Theorems 4.3–4.5 we can read off for which of the classical forms the statement of Theorem 4.7 is valid.

THEOREM 5.1. *Restricting attention to the classical weights (1–5) and (1–9), statement (4–2) holds for*

$$(f, g) = \begin{cases} (e^{-x^2/2}, e^{-x^2}), \\ (x^{(a-1)/2}e^{-x/2}, x^a e^{-x}) & \text{with } x > 0, \\ (x^{(a-1)/2}(1-x)^{(b-1)/2}, x^a(1-x)^b) & \text{with } 0 < x < 1, \\ ((1+x^2)^{-(\alpha+1)/2}, (1+x^2)^{-\alpha}); \end{cases} \quad (5-1)$$

the statement (4–3) holds for

$$(f, g) = \begin{cases} (e^{-x/2}, e^{-x}) & \text{with } x > 0, \\ ((1-x)^{(a-1)/2}, (1-x)^a) & \text{with } 0 < x < 1; \end{cases}$$

(4–4) is valid for the particular pair of Jacobi weights

$$(f, g) = (x^{(a-1)/2}, x^a) \quad \text{with } 0 < x < 1;$$

(4–5) is valid for the particular pair of Jacobi weights

$$(f, g) = (1, 1) \quad \text{with } 0 < x < 1;$$

and (4–6) is valid for the weights

$$((1+x^2)^{-(n+1)/2}, (1+x^2)^{-n}).$$

Because the weights in Theorem 5.1 occur in the matrix ensembles listed in the Introduction, the theorems of Section 4 imply interrelationships between the different ensembles.

THEOREM 5.2. *The following relations hold between the above matrix ensembles under decimation, for all $n > 0$:*

$$\begin{aligned} \text{even}(\text{GOE}_{2n+1}) &= \text{GSE}_n, \\ \text{even}(\text{Symm}(n; \mathbb{C})) &= \text{Anti}(n; \mathbb{C}), \\ \text{even}(\text{Mat}(2p+1, 2q+1; \mathbb{R})) &= \text{Mat}(p, q; \mathbb{H}), \\ \text{even}(\text{Beta}(2p_1+1, 2p_2+1, 2q+1; \mathbb{R})) &= \text{Beta}(p_1, p_2, q; \mathbb{H}), \\ \text{even}(\text{GOE}_n \cup \text{GOE}_{n+1}) &= \text{GUE}_n, \\ \text{even}(\text{Symm}(n; \mathbb{C}) \cup \text{Symm}(n; \mathbb{C})) &= \text{Mat}(n, n; \mathbb{C}), \\ \text{even}(\text{Symm}(n; \mathbb{C}) \cup \text{Symm}(n+1; \mathbb{C})) &= \text{Mat}(n+1, n; \mathbb{C}), \\ \text{even}(\text{Mat}(p, q; \mathbb{R}) \cup \text{Mat}(p+1, q+1; \mathbb{R})) &= \text{Mat}(p, q; \mathbb{C}), \\ \text{even}(\text{Beta}(p_1, p_2, q; \mathbb{R}) \cup \text{Beta}(p_1+1, p_2+1, q+1; \mathbb{R})) &= \text{Beta}(p_1, p_2, q; \mathbb{C}). \end{aligned}$$

REMARK 1. It would be very nice to have a direct, matrix-theoretic, proof of any of these relations.

REMARK 2. There are actually a few more relations, all of which follow from the preceding ones together with the relation

$$\text{Mat}(n+1, n; \mathbb{R}) = \text{Symm}(n; \mathbb{C}).$$

Again, a matrix-theoretic proof of this would be nice.

We now turn our attention to the implications of Theorem 5.1 with respect to gap probabilities. In circular ensemble theory the results (1–7) and (1–8) were shown [8] to imply interrelationships between the probability of an eigenvalue free region amongst the various symmetry classes. With $E^{(\beta)}(p; J; n)$ denoting the probability that, for the ensembles COE_n ($\beta = 1$), CUE_n ($\beta = 2$) and CSE_n ($\beta = 4$), there are exactly p eigenvalues in the interval J , the interrelationships are

$$\begin{aligned} E^{(2)}(0; (0, s); n) &= E^{(1)}(0; (0, s); n)(E^{(1)}(0; (0, s); n) + E^{(1)}(1; (0, s); n)), \\ E^{(4)}(p; (0, s); n) &= E^{(1)}(2p; (0, s); 2n) \\ &\quad + \frac{1}{2}E^{(1)}(2p-1; (0, s); 2n) + \frac{1}{2}E^{(1)}(2p+1; (0, s); 2n), \end{aligned} \quad (5-2)$$

where

$$E^{(\beta)}(p; J; n) := 0, \quad \text{for } p < 0. \quad (5-3)$$

Similar interrelationships between gap probabilities, but now with the eigenvalue free interval including an endpoint of the support of the interval, can be deduced from the pairs of statements of Theorem 4.7.

THEOREM 5.3. *Let $E^{(\beta)}(p; J; g; n)$ denote the probability that, for the ensembles $\text{OE}_n(g)$ ($\beta = 1$), $\text{UE}_n(g)$ ($\beta = 2$) and $\text{SE}_n(g)$ ($\beta = 4$) the interval J contains exactly n eigenvalues.*

(i) *The statements (4–2) imply*

$$\begin{aligned} E^{(2)}(0; J; g; 2n) &= E^{(1)}(0; J; f; 2n)E^{(1)}(0; J; f; 2n+1) \\ &\quad + E^{(1)}(0; J; f; 2n)E^{(1)}(1; J; f; 2n+1) + E^{(1)}(0; J; f; 2n+1)E^{(1)}(1; J; f; 2n), \\ E^{(4)}(p; J; (g/f)^2; n) &= E^{(1)}(2p; J; f; 2n+1) + E^{(1)}(2p+1; J; f; 2n+1), \end{aligned} \quad (5-4)$$

for $J = (-\infty, -s)$ or (s, ∞) .

(ii) *The statements (4–3) imply*

$$\begin{aligned} E^{(2)}(0; (s, \infty); g; 2n) &= (E^{(1)}(0; (s, \infty); f; 2n))^2 \\ &\quad + 2E^{(1)}(0; (s, \infty); f; 2n)E^{(1)}(1; (s, \infty); f; 2n), \\ E^{(4)}(p; (s, \infty); (g/f)^2; n) &= E^{(1)}(2p; (s, \infty); f; 2n) + E^{(1)}(2p+1; (s, \infty); f; 2n) \end{aligned} \quad (5-5)$$

and

$$\begin{aligned} E^{(2)}(0; (-\infty, -s); g; 2n) &= (E^{(1)}(0; (-\infty, -s); f; 2n))^2, \\ E^{(4)}(p; (-\infty, -s); (g/f)^2; n) &= E^{(1)}(2p; (-\infty, -s); f; 2n) \\ &\quad + E^{(1)}(2p-1; (-\infty, -s); f; 2n). \end{aligned} \quad (5-6)$$

(iii) *The statements (4-4) imply*

$$E^{(2)}(0; (s, \infty); g; 2n) = (E^{(1)}(0; (s, \infty); f; 2n))^2,$$

$$E^{(4)}(p; (s, \infty); (g/f)^2; n) = E^{(1)}(2p; (s, \infty); f; 2n) + E^{(1)}(2p-1; (s, \infty); f; 2n) \quad (5-7)$$

and

$$E^{(2)}(0; (-\infty, -s); g; 2n) = (E^{(1)}(0; (-\infty, -s); f; 2n))^2$$

$$+ 2E^{(1)}(0; (-\infty, -s); f; 2n)E^{(1)}(1; (-\infty, -s); f; 2n),$$

$$E^{(4)}(p; (-\infty, -s); (g/f)^2; n) = E^{(1)}(2p; (-\infty, -s); f; 2n)$$

$$+ E^{(1)}(2p+1; (-\infty, -s); f; 2n). \quad (5-8)$$

(iv) *The statements (4-5) imply*

$$E^{(2)}(0; J; g; 2n) = E^{(1)}(0; J; f; 2n)E^{(1)}(0; J; f; 2n-1),$$

$$E^{(4)}(p; J; (g/f)^2; n) = E^{(1)}(2p; J; f; 2n-1) + E^{(1)}(2p-1; J; f; 2n-1), \quad (5-9)$$

for $J = (-\infty, -s)$ or (s, ∞) .

(v) *The statements (4-6) imply the relations (5-2) with n replaced by $2n$ in the first equation and $(0, s)$ replaced throughout by J , $J = (-\infty, -s)$ or (s, ∞) .*

PROOF. We will consider only the deductions from (4-2), as the other cases are similar. Let J be a single interval which includes an endpoint of the support of f and g . From the first statement in (4-2) we see that the event of a sequence of eigenvalues from $\text{UE}_{2n}(g)$ not being contained in J occurs in three ways relative to the ensemble $\text{OE}_{2n}(f) \cup \text{OE}_{2n+1}(f)$: (i) the eigenvalues from $\text{OE}_{2n}(f)$ and those from $\text{OE}_{2n+1}(f)$ are not contained in J ; or (ii) one eigenvalue from $\text{OE}_{2n+1}(f)$ is contained in J and no eigenvalue from $\text{OE}_{2n}(f)$ is contained in J (note that the one eigenvalue must be either the largest (smallest) eigenvalue when J contains the right (left) hand end point); or (iii) one eigenvalue from $\text{OE}_{2n}(f)$ is contained in J and no eigenvalue from $\text{OE}_{2n+1}(f)$ is contained in J . This gives the first equation in (5-4). From the second statement in (4-2) we see that the event of a sequence of eigenvalues from $\text{SE}_n((g/f)^2)$ containing p eigenvalues in J can occur in two ways relative to $\text{OE}_{2n+1}(f)$: (i) there are $2p$ eigenvalues from $\text{OE}_{2n+1}(f)$ in J ; or (ii) there are $2p+1$ eigenvalues from $\text{OE}_{2n+1}(f)$ in J (of which $p+1$ are integrated out in forming $\text{even}(\text{OE}_{2n+1}(f))$). This implies the second equation in (5-4). \square

Recalling (5-3) we see that in the case $p = 0$ the equations (5-6), (5-7) and (5-9) give particularly simple interrelationships between the $E^{(\beta)}(0; \dots)$. In fact referring back to Theorem 5.1 for the permissible pairs (f, g) in these cases it is a simple exercise in changing variables to compute the $E^{(\beta)}(0; \dots)$ in terms of elementary functions. Recalling that

$$E^{(\beta)}(0; J; w; n) := \frac{1}{C} \int_{\bar{J}} dx_0 \cdots \int_{\bar{J}} dx_{n-1} \prod_{l=0}^{n-1} w(x_l) \prod_{0 \leq j < k \leq n-1} |x_k - x_j|^\beta,$$

where $\bar{J} = (-\infty, \infty) - J$ and C is such that $E^{(\beta)}(0; \emptyset; w; n) = 1$ we find

$$\begin{aligned} E^{(1)}(0; (0, s); e^{-x/2}; n) &= e^{-sn/2}, \\ E^{(2)}(0; (0, s); e^{-x}; n) &= E^{(4)}(0; (0, s); e^{-x}; n) = e^{-sn}, \\ E^{(1)}(0; (0, s); (1-x)^{(a-1)/2}; n) &= E^{(1)}(0; (1-s, 1); x^{(a-1)/2}; n) = (1-s)^{n(n+a)/2}, \\ E^{(2)}(0; (0, s); (1-x)^a; n) &= E^{(2)}(0; (1-s, 1); x^a; n) = (1-s)^{n(a+n)}, \\ E^{(4)}(0; (0, s); (1-x)^{a+1}; n) &= E^{(4)}(0; (1-s, 1); x^{a+1}; n) = (1-s)^{2n^2+na} \end{aligned}$$

(in the first two cases the weight functions are restricted to $x > 0$, while in the remaining cases $0 < x < 1$). The equations (5–6), (5–7) and (5–9) for $p = 0$ can be checked immediately.

The pairs of equations (5–4), (5–5) and (5–8) contain $E^{(1)}(1; \dots)$ as well as $E^{(\beta)}(0; \dots)$. In the equations (5–5) and (5–8) the dependence on $E^{(1)}(1; \dots)$ can be eliminated. Noting from Theorem 5.1 the allowed pairs (f, g) for the validity of equations (5–5) and (5–8) the following result is obtained.

PROPOSITION 5.4. *Assume that $(f, g) = (e^{-x/2}, e^{-x})$ with $x > 0$ and that $J = (s, \infty)$; or that $(f, g) = ((1-x)^{(a-1)/2}, (1-x)^a)$ with $0 < x < 1$ and that $J = (1-s, 1)$. Then*

$$E^{(4)}(0; J; (g/f)^2; n) = \frac{1}{2} \left(E^{(1)}(0; J; f; 2n) + \frac{E^{(2)}(0; J; g; 2n)}{E^{(1)}(0; J; f; 2n)} \right). \quad (5-10)$$

In the scaled $n \rightarrow \infty$ limit, as appropriate for the particular choice of weight function in (5–1), the pair of equations (4–2) also imply an equation of the form (5–10). First consider the Gaussian ensembles with the scaling [11]

$$x \mapsto (2n)^{1/2} + \frac{x}{2^{1/2}n^{1/6}},$$

which corresponds to studying the distribution of the eigenvalues at the (soft) edge of the leading order support of the spectrum. Defining

$$\begin{aligned} E_{\text{soft}}^{(1)}(p; (s, \infty)) &:= \lim_{n \rightarrow \infty} E^{(1)} \left(p; \left((n)^{1/2} + \frac{s}{2^{1/2}n^{1/6}}, \infty \right) e^{-x^2/2}; n \right) \\ E_{\text{soft}}^{(2)}(p; (s, \infty)) &:= \lim_{n \rightarrow \infty} E^{(2)} \left(p; \left((n)^{1/2} + \frac{s}{2^{1/2}n^{1/6}}, \infty \right) e^{-x^2}; n \right) \\ E_{\text{soft}}^{(4)}(p; (s, \infty)) &:= \lim_{n \rightarrow \infty} E^{(4)} \left(p; \left((n)^{1/2} + \frac{s}{2^{1/2}n^{1/6}}, \infty \right) e^{-x^2}; n/2 \right), \end{aligned}$$

(the existence of these limits is known from explicit calculation [27]; see below). The equations (5–4) imply:

PROPOSITION 5.5. *For the scaled infinite Gaussian ensembles at the soft edge*

$$E_{\text{soft}}^{(4)}(0; (s, \infty)) = \frac{1}{2} \left(E_{\text{soft}}^{(1)}(0; (s, \infty)) + \frac{E_{\text{soft}}^{(2)}(0; (s, \infty))}{E_{\text{soft}}^{(1)}(0; (s, \infty))} \right), \quad (5-11)$$

$$E_{\text{soft}}^{(1)}(1; (s, \infty)) = E_{\text{soft}}^{(4)}(0; (s, \infty)) - E_{\text{soft}}^{(1)}(0; (s, \infty)). \tag{5-12}$$

As alluded to above, the $E_{\text{soft}}^{(\beta)}(0; (s, \infty))$ are known exactly from the work of Tracy and Widom [27]. To present these results, let $q(s)$ denote the solution of the particular Painlevé II equation

$$q'' = sq + 2q^3,$$

which satisfies the boundary condition $q(s) \sim \text{Ai}(s)$ as $s \rightarrow \infty$. Then we have

$$\begin{aligned} E_{\text{soft}}^{(2)}(0; (s, \infty)) &= \exp\left(-\int_s^\infty (t-s)q^2(t) dt\right), \\ (E_{\text{soft}}^{(1)}(0; (s, \infty)))^2 &= E_{\text{soft}}^{(2)}(0; (s, \infty)) \exp\left(-\int_s^\infty q(t) dt\right), \\ (E_{\text{soft}}^{(4)}(0; (s, \infty)))^2 &= E_{\text{soft}}^{(2)}(0; (s, \infty)) \cosh^2\left(\frac{1}{2}\int_s^\infty q(t) dt\right) \end{aligned} \tag{5-13}$$

(in [27] $E_{\text{soft}}^{(4)}$ is defined with $s \mapsto s/2^{1/2}$ relative to our definition). Equation (5-11) is immediately seen to be satisfied, while the second equation gives

$$(E_{\text{soft}}^{(1)}(1; (s, \infty)))^2 = E_{\text{soft}}^{(2)}(0; (s, \infty)) \sinh^2\left(\frac{1}{2}\int_s^\infty q(t) dt\right). \tag{5-14}$$

Next consider the scaled limit at an edge for which the weight function is strictly zero on one side. For the classical ensembles this occurs in the Laguerre and Jacobi case; for definiteness consider the Laguerre case. The appropriate scaling is [11]

$$x \mapsto \frac{x}{4n},$$

and we define

$$\begin{aligned} E_{\text{hard}}^{(1)}(p; (0, s); (a-1)/2) &:= \lim_{n \rightarrow \infty} E^{(1)}(p; (0, s/4n); x^{(a-1)/2}e^{-x}; n), \\ E_{\text{hard}}^{(2)}(p; (0, s); a) &:= \lim_{n \rightarrow \infty} E^{(2)}(p; (0, s/4n); x^a e^{-x}; n), \\ E_{\text{hard}}^{(4)}(p; (0, s); a+1) &:= \lim_{n \rightarrow \infty} E^{(4)}(p; (0, s/4n); x^{a+1}e^{-x}; n/2) \end{aligned}$$

(the existence of these limits for general $a > -1$ can be deduced from the existence of the k -point distributions in the same scaled limits [22]). Use of (4-2) then gives the analogue of Proposition 5.5 for the hard edge.

PROPOSITION 5.6. *For the scaled infinite Laguerre ensembles at the hard edge, we have*

$$E_{\text{hard}}^{(4)}(0; (0, s); a+1) = \frac{1}{2}\left(E_{\text{hard}}^{(1)}(0; (0, s); \frac{1}{2}(a-1)) + \frac{E_{\text{hard}}^{(2)}(0; (0, s); a)}{E_{\text{hard}}^{(1)}(0; (0, s); (a-1)/2)}\right) \tag{5-15}$$

and

$$E_{\text{hard}}^{(1)}(1; (0, s); \frac{1}{2}(a-1)) = E_{\text{hard}}^{(4)}(0; (0, s); a+1) - E_{\text{hard}}^{(1)}(0; (0, s); (a-1)/2).$$

Exact Pfaffian formulas are known for

$$E_{\text{hard}}^{(1)}(0; (0, s); (a-1)/2) \quad \text{and} \quad E_{\text{hard}}^{(4)}(0; (0, s); a+1)$$

in the case of a an odd positive integer [23], while $E_{\text{hard}}^{(2)}(0; (0, s); a)$ can then be expressed as a determinant [13] (the dimension of the Pfaffians and the determinants are proportional to a), although (5–15) is not a natural consequence of these formulas. There are also multiple integral expressions for the same expression [10], but again they do not naturally satisfy (5–15).

Note Added. An analogous formula to that in (5–13) for $E_{\text{soft}}^{(1)}(0; (s, \infty))$, but now involving a certain Painlevé V transcendent, has recently been derived [12] for

$$E_{\text{hard}}^{(1)}(0; (0, s); \frac{1}{2}(a-1)),$$

and formula (5–15) then used to compute $E_{\text{hard}}^{(4)}(0; (0, s); a+1)$.

6. Distribution Functions for Superimposed Spectra

In general, for a symmetric PDF $p(x_0, \dots, x_{n-1})$ the k -point distribution function ρ_k is defined by

$$\rho_k(x_0, \dots, x_{k-1}) := n(n-1) \cdots (n-k+1) \int_{(-\infty, \infty)^{n-k}} p(x_0, \dots, x_{n-1}) dx_k \cdots dx_{n-1}. \quad (6-1)$$

In this section we take up the task of computing ρ_k for $\text{even}(M)$, $\text{odd}(M)$, $\text{alt}(M)$ with $M = \text{OE}_n(f) \cup \text{OE}_n(f)$ and $\text{even}(M)$, $\text{odd}(M)$ with $M = \text{OE}_n(f) \cup \text{OE}_{n+1}(f)$.

For $M = \text{OE}_n(f) \cup \text{OE}_n(f)$, write

$$\begin{aligned} D^{\text{even}(M)}(x_0, \dots, x_{n-1}) &:= \det(F_j(x_i) - F_j(I))_{0 \leq i, j < n}, \\ D^{\text{odd}(M)}(x_0, \dots, x_{n-1}) &:= \det(F_j(x_i))_{0 \leq i, j < n}, \\ D^{\text{alt}(M)}(x_0, \dots, x_{n-1}) &:= \det(F_j(x_i) - \frac{1}{2}F_j(I))_{0 \leq i, j < n} \quad (n \text{ even}), \\ D^{\text{alt}(M)}(x_0, \dots, x_{n-1}) &:= \det \begin{pmatrix} 0 & (F_j(I))_{0 \leq j < n} \\ (1)_{1 \leq i < n-1} & (F_j(x_i))_{0 \leq i, j < n} \end{pmatrix} \quad (n \text{ odd}), \end{aligned} \quad (6-2)$$

and for $M = \text{OE}_n(f) \cup \text{OE}_{n+1}(f)$ let

$$\begin{aligned} D^{\text{even}(M)}(x_0, \dots, x_{n-1}) &:= \det \begin{pmatrix} (F_j(x_i))_{0 \leq i < n, 0 \leq j < n+1} \\ (F_j(I))_{0 \leq j < n+1} \end{pmatrix}, \\ D^{\text{odd}(M)}(x_0, \dots, x_{n-1}) &:= \det \begin{pmatrix} (1)_{0 \leq j < n+1} \\ (F_j(x_i))_{0 \leq i < n, 0 \leq j < n+1} \end{pmatrix}. \end{aligned} \quad (6-3)$$

In each case, workings contained in the proofs of Theorems 4.2 and 4.6 show (after relabelling the coordinates) that the PDF is proportional to

$$\prod_{i=0}^{n-1} f(x_i) \Delta(x_0, \dots, x_{n-1}) D(x_0, \dots, x_{n-1}) \quad (6-4)$$

for D as specified. Now introduce a set of functions $\{\eta_j(x)\}_{0 \leq j < n}$ such that

$$D(x_0, \dots, x_{n-1}) \propto \det(\eta_j(x_i))_{0 \leq i, j < n}$$

and a set of monic polynomials $\{q_j(x)\}_{0 \leq j < n}$, q_j of degree j , such that the biorthogonality property

$$\int_{-\infty}^{\infty} f(x) q_i(x) \eta_j(x) dx = \delta_{i,j}$$

holds (assuming such biorthogonal families exist). The k -point distribution can be expressed in terms of these functions.

LEMMA 6.1. *For the PDF (6-4) and $\{\eta_j(x)\}_{0 \leq j < n}$, $\{q_j(x)\}_{0 \leq j < n}$ specified as above, we have*

$$\rho_k(x_0, \dots, x_{k-1}) = \prod_{j=0}^{k-1} f(x_j) \det \left(\sum_{l=0}^{n-1} q_l(x_i) \eta_l(x_j) \right)_{0 \leq i, j < k}. \quad (6-5)$$

PROOF. From the definitions of $\{\eta_j(x)\}_{0 \leq j < n}$ and $\{q_j(x)\}_{0 \leq j < n}$ we see that (6-4) is proportional to

$$\prod_{i=0}^{n-1} f(x_i) \det(q_j(x_i))_{0 \leq i, j < n} \det(\eta_j(x_i))_{0 \leq i, j < n}.$$

The biorthogonal property allows the integrations required by the definition (6-1) to be computed to give (6-5). \square

REMARK. Suppose for some $\{\xi_j(x)\}_{j=0, \dots, n-1}$ we can write

$$D(x_0, \dots, x_{n-1}) \propto \det(\xi_j(x_i))_{0 \leq i, j < n-1}.$$

It is easy to show [18; 6] that sufficient conditions for the existence of the biorthogonal sets is that

$$\det \left(\int_{-\infty}^{\infty} f(x) x^i \xi_j(x) dx \right)_{0 \leq i, j < p} \neq 0$$

for $p = 0, \dots, n-1$.

For f a classical weight and so of the form (1–5) or (1–9), the biorthogonal functions can be computed explicitly. This is possible because of the following special property of the classical weights and their corresponding orthogonality [1].

LEMMA 6.2. *Consider the pairs (f, g) of classical weight functions (5–1). Let $\{p_j(x)\}_{j=0,1,\dots}$ be the set of monic orthogonal polynomials, $p_j(x)$ of degree j corresponding to the weight function g , let $(p_k, p_k)_2$ denote their normalization with respect to integration over the measure $g(x)dx$, and define γ_k so that $\gamma_k(p_k, p_k)_2$ equals respectively $1, \frac{1}{2}, \frac{1}{2}(2k+2+a+b)$, or $\alpha-k-1$ in the four cases. With*

$$\mathbf{n} := \frac{1}{f(x)} \frac{d}{dx} \frac{g(x)}{f(x)}, \quad c_k := \gamma_k(p_k, p_k)_2 (p_{k+1}, p_{k+1})_2$$

we have

$$\mathbf{n} p_k(x) = -\frac{c_k}{(p_{k+1}, p_{k+1})_2} p_{k+1}(x) + \frac{c_{k-1}}{(p_{k-1}, p_{k-1})_2} p_{k-1}(x).$$

PROOF. This is a simple consequence of the following property [2]:

$$(\phi, \mathbf{n} \psi)_2 = -(\mathbf{n} \phi, \psi)_2. \quad \square$$

As a consequence, the determinant formulas (6–2) and (6–3) for $D(x_0, \dots, x_{n-1})$ can be simplified. Let

$$(u(x))_j := \sum_{l=j}^{\infty} \frac{(u(x), p_l(x))_2}{(p_l, p_l)_2} p_l(x) = u(x) - \sum_{l=0}^{j-1} \frac{(u(x), p_l(x))_2}{(p_l, p_l)_2} p_l(x),$$

and define

$$\begin{aligned} r_j^{(1)}(x) &= \left(\frac{f(x)}{g(x)} \int_x^{\infty} f(t) dt \right)_j, \\ r_j^{(2)}(x) &= \left(\frac{f(x)}{g(x)} \int_{-\infty}^x f(t) dt \right)_j, \\ r_j^{(3)}(x) &= \left(\frac{f(x)}{g(x)} \left(\int_{-\infty}^x f(t) dt - \frac{1}{2} \int_{-\infty}^{\infty} f(t) dt \right) \right)_j, \\ r_j^{(4)}(x) &= \left(\frac{f(x)}{g(x)} \right)_j, \\ r_j^{(5)}(x) &= r_{j-1}^{(4)}(x) - \frac{(r_{j-1}^{(4)}, r_{j-1}^{(4)})_2}{(r_{j-1}^{(4)}, r_{j-1}^{(2)})_2} r_{j-1}^{(2)}(x). \end{aligned}$$

Then, by adding appropriate linear combinations of the columns in the determinant formulas (6–2) and (6–3) and making use of Lemma 6.2, we readily find for

$$M = \text{OE}_n(f) \cup \text{OE}_n(f)$$

$$D^{\text{even}(M)}(x_0, \dots, x_{n-1}) \propto \prod_{i=1}^{n-1} \frac{g(x_i)}{f(x_i)} \det \left((p_j(x_i))_{\substack{0 \leq i < n \\ 0 \leq j < n-1}} (r_{n-1}^{(1)}(x_i))_{0 \leq i < n} \right),$$

$$D^{\text{odd}(M)}(x_0, \dots, x_{n-1}) \propto \prod_{i=1}^{n-1} \frac{g(x_i)}{f(x_i)} \det \left((p_j(x_i))_{\substack{0 \leq i < n \\ 0 \leq j < n-1}} (r_{n-1}^{(2)}(x_i))_{0 \leq i < n} \right),$$

$$D^{\text{alt}(M)}(x_0, \dots, x_{n-1}) \propto \prod_{i=1}^{n-1} \frac{g(x_i)}{f(x_i)} \det \left((p_j(x_i))_{\substack{0 \leq i < n \\ 0 \leq j < n-1}} (r_{n-1}^{(3)}(x_i))_{0 \leq i < n} \right) \\ (n \text{ even}),$$

$$D^{\text{alt}(M)}(x_0, \dots, x_{n-1}) \propto \prod_{i=1}^{n-1} \frac{g(x_i)}{f(x_i)} \det \left((p_j(x_i))_{\substack{0 \leq i < n \\ 0 \leq j < n-2}} (r_{n-1}^{(4)}(x_i))_{0 \leq i < n} \right) \\ (n \text{ odd}),$$

while for $M = \text{OE}_n(f) \cup \text{OE}_{n+1}(f)$

$$D^{\text{even}(M)}(x_0, \dots, x_{n-1}) \propto \prod_{i=1}^{n-1} \frac{g(x_i)}{f(x_i)} \det \left(p_j(x_i) \right)_{0 \leq i, j < n},$$

$$D^{\text{odd}(M)}(x_0, \dots, x_{n-1}) \\ \propto \prod_{i=1}^{n-1} \frac{g(x_i)}{f(x_i)} \det \left((p_j(x_i))_{\substack{0 \leq i < n \\ 0 \leq j < n-2}} (r_{n-2}^{(4)}(x_i))_{0 \leq i < n} (r_{n-1}^{(5)}(x_i))_{0 \leq i < n} \right).$$

In each case, setting $\eta_j(x) / \int_{-\infty}^{\infty} f(x) p_i(x) \eta_j(x) dx$ equal to $g(x)/f(x)$ times the function in column j and $q_i(x) = p_i(x)$ we have

$$\int_{-\infty}^{\infty} f(x) q_i(x) \eta_j(x) dx = \delta_{i,j} \quad (i, j = 0, \dots, n-1),$$

which is the desired biorthogonality property. Hence substitution of these values into (6-5) gives the k -point distribution in each case.

In particular with $M = \text{OE}_n(f) \cup \text{OE}_{n+1}(f)$ and f one of the classical weights in (5-1) we read off that

$$\rho_k^{\text{even}(M)}(x_0, \dots, x_{k-1}) = \det \left((g(x_i)g(x_j))^{1/2} \sum_{l=0}^{n-1} \frac{p_l(x_i)p_l(x_j)}{(p_l, p_l)_2} \right)_{0 \leq i, j < k}.$$

This is the well known expression for ρ_k in $\text{UE}(g)$, and thus is in keeping with the result of Theorem 4.3, giving $\text{even}(\text{OE}_n(f) \cup \text{OE}_{n+1}(f)) = \text{UE}_n(g)$ for each of the pairs (f, g) in (5-1). Furthermore, the Christoffel-Darboux formula evaluates the sum as

$$S_2(x, y) := (g(x)g(y))^{1/2} \sum_{l=0}^{n-1} \frac{p_l(x)p_l(y)}{(p_l, p_l)_2} \\ = \frac{(g(x)g(y))^{1/2} p_n(x)p_{n-1}(y) - p_{n-1}(x)p_n(y)}{(p_{n-1}, p_{n-1})_2 x - y}. \quad (6-6)$$

7. Distribution Functions for Alternate Eigenvalues in a Single OE_n

The k -point distribution function for the alternate eigenvalues in a single OE_n has a different structure to ρ_k for the superimposed OE_n spectra. The cases n even and n odd must be treated separately.

Case of n even. Consider first $\text{even}(\text{OE}_n(f))$ with n even. From the manipulations sketched in the proof of Theorem 4.7 we have that the PDF of this ensemble is given by

$$\frac{1}{C} \prod_{l=0}^{n/2-1} f(x_{2l}) \det \left(\int_{x_{2i}}^{\infty} t^j f(t) dt \right)_{\substack{0 \leq i < n \\ 0 \leq j < 2n}}. \quad (7-1)$$

To perform the integration required by (6-1) we introduce the skew inner product

$$\begin{aligned} \langle u|v \rangle_1 &:= \frac{1}{2} \int_{-\infty}^{\infty} dx f(x) \left(u(x) \int_x^{\infty} dy f(y)v(y) - v(x) \int_x^{\infty} dy f(y)u(y) \right) \\ &= \frac{1}{2} \int_{-\infty}^{\infty} dx f(x)u(x) \int_{-\infty}^{\infty} dy v(y) \text{sgn}(y-x), \end{aligned} \quad (7-2)$$

together with a corresponding family of monic skew orthogonal polynomials $\{R_i(x)\}_{i=0,1,\dots}$ which are defined so that

$$\begin{aligned} \langle R_{2i}|R_{2j+1} \rangle_1 &= -\langle R_{2j+1}|R_{2i} \rangle_1 = r_j \delta_{i,j}, \\ \langle R_{2i}|R_{2j} \rangle_1 &= \langle R_{2i+1}|R_{2j+1} \rangle_1 = 0. \end{aligned} \quad (7-3)$$

Note that the skew orthogonality property still holds if we make the replacement

$$R_{2i+1}(x) \mapsto R_{2i+1}(x) + \gamma_{2i} R_{2i}(x)$$

for arbitrary γ_{2i} . However, a construction of Gram-Schmidt type shows that $\{R_i(x)\}_{i=0,1,\dots}$ is unique up to this transformation.

We will start by expressing (7-1) as a quaternion determinant involving $\{R_i(x)\}_{i=0,1,\dots}$ and then show how the property (7-3) can be used to perform the integrations. This requires the definition of a quaternion determinant. We regard a quaternion as a 2×2 matrix, and a quaternion matrix as a matrix with quaternion elements. With n even and

$$Z_n := \mathbf{1}_{n/2} \otimes \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

a $n/2 \times n/2$ quaternion matrix Q is said to be self dual if

$$Q^D := Z_n Q^T Z_n^{-1} = Q.$$

In terms of its 2×2 sub-blocks this means that the quaternion element in position (kj) is related to the element in position (jk) , $j < k$ by

$$q_{kj} = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \quad \text{if} \quad q_{jk} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

Now for a self dual quaternion matrix the determinant, to be denoted qdet , is defined by [9]

$$\text{qdet } Q = \sum_{P \in S_{n/2}} (-1)^{n/2-l} \prod_l (q_{ab} q_{bc} \cdots q_{da})^{(0)} \quad (7-4)$$

where the superscript (0) denotes the operation $\frac{1}{2}\text{Tr}$, P is any permutation of the indices $(1, \dots, n/2)$ consisting of l exclusive cycles of the form $(a \rightarrow b \rightarrow c \rightarrow \cdots d \rightarrow a)$ and $(-1)^{n/2-l}$ is the parity of P . Furthermore, $\text{qdet } Q$ is related to the Pfaffian via the formula [9]

$$\text{qdet } Q = \text{Pf } QZ_n^{-1},$$

which since $(\text{Pf } QZ_n^{-1})^2 = \det Q$ (where Q is regarded as an ordinary $n \times n$ matrix) implies [17]

$$\det Q = \text{qdet}(QQ^D), \quad (7-5)$$

assuming $\det Q$ is positive.

PROPOSITION 7.1. *With $p(x_0, x_2, \dots, x_{n-2})$ denoting the probability density function (7-1), $\{R_i(x)\}_{i=0,1,\dots}$ the monic orthogonal polynomials with respect to (7-2) and $\{r_i\}_{i=0,1,\dots}$ the corresponding normalizations we can write*

$$p(x_0, x_2, \dots, x_{n-2}) = \frac{1}{C} \prod_{k=0}^{n/2-1} (2r_k) \text{qdet}(T(x_{2j}, x_{2k}))_{0 \leq j, k < n/2}, \quad (7-6)$$

where

$$\begin{aligned} T(x, y) &:= \sum_{k=0}^{n/2-1} \frac{1}{2r_k} (\chi_k(y) \chi_k^D(x))^T = \begin{pmatrix} S(x, y) & I(x, y) \\ D(x, y) & S(y, x) \end{pmatrix}, \\ \chi_k(x) &:= \begin{pmatrix} f(x) R_{2k}(x) & f(x) R_{2k+1}(x) \\ \int_x^\infty f(t) R_{2k}(t) dt & \int_x^\infty f(t) R_{2k+1}(t) dt \end{pmatrix}, \\ S(x, y) &= \sum_{k=0}^{N/2-1} \frac{f(y)}{2r_k} \left(R_{2k}(y) \int_x^\infty f(t) R_{2k+1}(t) dt - R_{2k+1}(y) \int_x^\infty f(t) R_{2k}(t) dt \right), \\ I(x, y) &= - \int_x^y S(x, y') dy', \\ D(x, y) &= \frac{\partial}{\partial y} S(x, y). \end{aligned} \quad (7-7)$$

PROOF. Because the polynomials $\{R_k(x)\}_{k=0,1,\dots}$ are monic we can add multiples of columns in (7-1) to obtain

$$p(x_0, x_2, \dots, x_{n-2}) = \frac{1}{C} \det \left(\begin{array}{c} f(x_{2j})R_{k-1}(x_{2j}) \\ \int_{x_{2j}}^\infty f(t)R_k(t) dt \end{array} \right)_{\substack{0 \leq j < n/2 \\ 0 \leq k < n}} = \frac{1}{C} \prod_{k=0}^{n/2-1} (2r_k) \\ \times \det \left(\begin{array}{cc} f(x_{2j})(2r_k)^{-1/2}R_{2k}(x_{2j}) & f(x_{2j})(2r_k)^{-1/2}R_{2k+1}(x_{2j}) \\ (2r_k)^{-1/2} \int_{x_{2j}}^\infty f(t)R_{2k}(t) dt & (2r_k)^{-1/2} \int_{x_{2j}}^\infty f(t)R_{2k+1}(t) dt \end{array} \right)_{0 \leq j, k < n/2} \quad (7-8)$$

Application of (7-5) and the formula $\text{qdet } A = \text{qdet } A^T$ gives the formula (7-6) with $S(x, y)$ as specified and formulas for $I(x, y)$ and $D(x, y)$ which are easily seen to be expressible in terms of $S(x, y)$ as stated. \square

A special feature of $T(x, y)$, which follows from its definition in (7-7) in terms of $\chi_k(y)\chi_k^D(x)$ and the skew orthogonality of $\{R_k(x)\}_{k=0,1,\dots}$ with respect to (7-2), is the integration formulas

$$\int_{-\infty}^\infty T(x, x) dx = N/2, \\ \int_{-\infty}^\infty T(x, y)T(y, z) dy = T(x, z). \quad (7-9)$$

As a consequence of (7-9) and the quaternion formula (7-6), the integrations required to compute (6-1) can be carried out. Thus with (7-9) holding it is generally true that [17]

$$\int_{-\infty}^\infty dx_{2m} \text{qdet} (T(x_{2i}, x_{2j}))_{0 \leq i, j \leq m} = (n/2 - (m-1)) \text{qdet} (T(x_{2i}, x_{2j}))_{0 \leq i, j \leq m-1}. \quad (7-10)$$

Consequently we see from (7-6) that

$$\rho_k(x_0, \dots, x_{2k-2}) = \text{qdet} (T(x_{2i}, x_{2j}))_{0 \leq i, j < k}. \quad (7-11)$$

If instead of considering $\text{even}(\text{OE}_n(f))$ we consider $\text{odd}(\text{OE}_n(f))$, the reasoning is essentially unchanged. Thus (7-6) and (7-7) hold with the replacements

$$\int_x^\infty \mapsto \int_{-\infty}^x \quad \text{and} \quad \{x_0, x_2, \dots, x_{n-2}\} \mapsto \{x_1, x_3, \dots, x_{n-1}\}, \quad (7-12)$$

and with this modification of $T(x, y)$ the formula (7-11) for ρ_k holds with the replacements

$$\{x_0, x_2, \dots, x_{2k-2}\} \mapsto \{x_1, \dots, x_{2k-1}\}. \quad (7-13)$$

The structure of (7-11) with $T(x, y)$ given by (7-7) is very similar to the general expression for ρ_k as computed for the ensemble $\text{SE}_n((g/f)^2)$. First it is

necessary to introduce monic skew orthogonal polynomials $\{Q_k(x)\}_{k=0,1,\dots}$ and corresponding normalizations $\{q_k\}_{k=0,1,\dots}$ with respect to the skew inner product

$$\langle u|v\rangle_4 := \int_{-\infty}^{\infty} dx (g(x)/f(x))^2 (u(x)v'(x) - u'(x)v(x)).$$

We then have [24] (see also [28])

$$\rho_k(x_0, \dots, x_{k-1}) = \text{qdet} (T_4(x_i, x_j))_{0 \leq i, j \leq k} \quad (7-14)$$

where

$$\begin{aligned} T_4(x, y) &:= \begin{pmatrix} S_4(x, y) & I_4(x, y) \\ D_4(x, y) & S_4(y, x) \end{pmatrix}, \\ S_4(x, y) &= \sum_{k=0}^{N-1} \frac{f(y)}{2q_k} \left(Q_{2k}(y) \frac{d}{dx} (f(x)Q_{2k+1}(x)) - Q_{2k+1}(x) \frac{d}{dx} (f(x)Q_{2k}(x)) \right), \\ I_4(x, y) &= - \int_x^y S_4(x, y') dy', \\ D_4(x, y) &= \frac{\partial}{\partial y} S_4(x, y). \end{aligned} \quad (7-15)$$

Case of n odd. The PDF for the distribution $\text{even}(\text{OE}_n(f))$ with n odd is

$$\frac{1}{C} \prod_{l=0}^{(n-3)/2} f(x_{2l}) \det \begin{pmatrix} \begin{pmatrix} x_{2i}^{2j} \\ \int_{x_{2j}}^{\infty} f(t)t^k dt \end{pmatrix}_{\substack{0 \leq i < (n-1)/2 \\ 0 \leq j < n}} \\ \left(\int_{-\infty}^{\infty} w_1(t)t^j dt \right)_{0 \leq j < n} \end{pmatrix}.$$

As in (7-8), we can introduce the monic polynomials $\{R_j(x)\}_{j=0,1,\dots}$ to rewrite this as

$$\frac{1}{C} \det \begin{pmatrix} \begin{pmatrix} f(x_{2i})R_j(x_{2i}) \\ \int_{x_{2i}}^{\infty} f(t)R_j(t) dt \end{pmatrix}_{\substack{0 \leq i < (n-1)/2 \\ 0 \leq j < n}} \\ \left(\int_{-\infty}^{\infty} f(t)R_j(t) dt \right)_{j=0, \dots, n-2} \end{pmatrix}.$$

Subtracting appropriate multiples of the last column from the columns $0, 1, \dots, n-2$ so as to eliminate the element of the column in the final row then gives

$$\frac{1}{C} \left(\int_{-\infty}^{\infty} f(t)R_{n-1}(t) dt \right) \det \begin{pmatrix} f(x_{2i})\hat{R}_j(x_{2i}) \\ \int_{x_{2i}}^{\infty} f(t)\hat{R}_j(t) dt \end{pmatrix}_{\substack{0 \leq i < (n-1)/2 \\ 0 \leq j < n}}, \quad (7-16)$$

where

$$\hat{R}_j(x) := R_j(x) - \left(\frac{\int_{-\infty}^{\infty} f(t)R_j(t) dt}{\int_{-\infty}^{\infty} f(t)R_{n-1}(t) dt} \right) R_{n-1}(x). \quad (7-17)$$

The determinant in (7-16) is formally the same as that in (7-8). Thus in the case n odd $p(x_0, x_2, \dots, x_{n-3})$ can be written as in (7-6) but with

$$n \mapsto n-1, \quad R_i \mapsto \hat{R}_i \quad (7-18)$$

and $C \mapsto C'$ for some normalization C' .

Now we can check from the definition (7-17) that for $j = 1, \dots, n - 1$ the polynomials \hat{R}_{j-1} satisfy the skew orthogonality property (7-3). This means that the integration formula (7-10) again applies in this modified setting and consequently the k -point distribution is given by

$$\rho_k(x_0, \dots, x_{2k-2}) = \text{qdet} \left(T^{\text{odd}}(x_{2i}, x_{2j}) \right)_{0 \leq i, j < k}, \tag{7-19}$$

where T^{odd} is defined as $T(x, y)$ in (7-7) but with the replacements (7-18). In the case of $\text{odd}(\text{OE}_n(f))$ the replacements (7-12) and (7-18) must be made in (7-11) and (7-7), and the replacement (7-13) made in (7-19).

7.1. Summation Formulas. It has already been remarked that ρ_k for $\text{SE}_n((g/f)^2)$ has the quaternion determinant form (7-14) and (7-15). Furthermore it is known [1] that with f one of the classical weights in (5-1), the quantity S_4 in (7-15) can be summed to give an expression independent of the skew orthogonal polynomials associated with g , and dependent only on the monic orthogonal polynomials $\{p_i(x)\}_{i=0,1,\dots}$ associated with the weight function $g(x)$. Explicitly,

$$2S_4(x, y) = \left(\frac{g(x)}{g(y)} \right)^{1/2} \frac{f(y)}{f(x)} S_2(x, y) \Big|_{n \mapsto 2n} - \gamma_{2n-1} f(y) p_{2n}(y) \int_x^\infty f(t) p_{2n-1}(t) dt, \tag{7-20}$$

where S_2 is specified by (6-6) and γ_{2n-1} by Lemma 6.2. Here we will use results from [1] to obtain an analogous summation for the quantity $S(x, y)$ in (7-7).

Suppose n is even and write

$$\Phi_j(x) := \frac{1}{2} \int_{-\infty}^\infty f(t) \text{sgn}(x - t) R_j(t) dt.$$

Then straightforward manipulation of the definition of $S(x, y)$ allows it to be rewritten

$$S(x, y) = \frac{1}{2} \left(S_1(x, y) - S_1(\infty, y) \right) \tag{7-21}$$

where

$$S_1(x, y) = \sum_{k=0}^{n/2-1} \frac{f(y)}{r_k} \left(\Phi_{2k}(x) R_{2k+1}(y) - \Phi_{2k+1}(x) R_{2k}(y) \right)$$

The quantity $S_1(x, y)$ occurs in the quaternion determinant formula for k -point distribution of $\text{OE}_n(f)$. With f one of the classical forms (5-1) it can be summed to give [1]

$$S_1(x, y) = \left(\frac{g(x)}{g(y)} \right)^{1/2} \frac{f(y)}{f(x)} S_2(x, y) \Big|_{n \mapsto n-1} + \gamma_{n-2} f(y) p_{n-1}(y) \frac{1}{2} \int_{-\infty}^\infty \text{sgn}(x - t) f(t) p_{n-2}(t) dt. \tag{7-22}$$

From this it follows that

$$S_1(\infty, y) = \gamma_{n-2} f(y) p_{n-1}(y) \frac{1}{2} \int_{-\infty}^{\infty} f(t) p_{n-2}(t) dt,$$

and so by (7-21) we can evaluate $S(x, y)$.

PROPOSITION 7.2. For (f, g) a classical pair (5-1), $\{p_j(x)\}_{j=0,1,\dots}$ monic orthogonal polynomials with respect to the weight function $g(x)$, and n even the quantity $S(x, y)$ in (7-7) has the evaluation

$$2S(x, y) = \left(\frac{g(x)}{g(y)} \right)^{1/2} \frac{f(y)}{f(x)} S_2(x, y) \Big|_{n \mapsto n-1} - \gamma_{n-2} f(y) p_{n-1}(y) \int_x^{\infty} f(t) p_{n-2}(t) dt \tag{7-23}$$

(compare (7-20)).

This summation fully determines $\text{even}(\text{OE}_n(f))$ with n even. For $\text{odd}(\text{OE}_n(f))$, n even, the prescription (7-12) says the replacement $\int_x^{\infty} \mapsto \int_{-\infty}^x$ should be made in (7-23).

It remains to consider the case n odd. Consider first $\text{even}(\text{OE}_n(f))$. In fact the formulas in [1] giving the analogous formula to (7-22) for n odd allows us to deduce that the summation (7-23) remains valid for n odd. For n odd comparison of (7-23) and (7-20) shows that

$$S(x, y) = S_4(x, y) \Big|_{n \mapsto (n-1)/2},$$

which because of the formulas (7-7) (with $n \mapsto n-1$), (7-11), (7-14), and (7-15) implies

$$\rho_k^{\text{even}(\text{OE}_{2n+1}(f))}(x_0, x_2, \dots, x_{2k-2}) = \rho_k^{\text{SE}_n((g/f)^2)}(x_0, x_2, \dots, x_{2k-2}).$$

This is equivalent to the second statement of (4-2), which we already know from Theorem 5.1 is valid for the pairs (f, g) in (5-1). In the case of $\text{odd}(\text{OE}_n(f))$ with n odd, again the prescription (7-12) says we simply make the replacement $\int_x^{\infty} \mapsto \int_{-\infty}^x$ in (7-23).

Acknowledgements

Forrester thanks J. Baik for drawing his attention to the conjecture noted in the paragraph above the paragraph containing (1-6), and acknowledges the Australian Research Council for financial support.

References

- [1] M. Adler, P. J. Forrester, T. Nagao, and P. van Moerbeke, "Classical skew orthogonal polynomials and random matrices", *J. Stat. Phys.* **99** (2000), 141-170.
- [2] M. Adler and P. van Moerbeke, "Matrix integrals, Toda symmetries, Virasoro constraints and orthogonal polynomials", *Duke Math. Journal* **80** (1995), 863-911.

- [3] A. Altland and M. R. Zirnbauer, “Nonstandard symmetry classes in mesoscopic normal-superconducting hybrid compounds”, *Phys. Rev. B* **55** (1997), 1142.
- [4] J. Baik and E. M. Rains, “Algebraic aspects of increasing subsequences”, preprint, math.CO/9905083, 1999.
- [5] J. Baik and E. M. Rains, “The asymptotics of monotone subsequences of involutions”, Preprint, math.CO/9905084, 1999.
- [6] A. Borodin, “Biorthogonal ensembles”, *Nucl. Phys. B* **536** (1999), 704–732.
- [7] F. J. Dyson, “Statistical theory of energy levels of complex systems I”, *J. Math. Phys.* **3** (1962), 140–156.
- [8] F. J. Dyson, “The three fold way. Algebraic structure of symmetry groups and ensembles in quantum mechanics”, *J. Math. Phys.* **3** (1962), 1199–1215.
- [9] F. J. Dyson, “Correlations between the eigenvalues of a random matrix”, *Commun. Math. Phys.* **19** (1970), 235–250.
- [10] P. J. Forrester, “Exact results and universal asymptotics in the Laguerre random matrix ensemble”, *J. Math. Phys.* **35** (1993), 2539–2551.
- [11] P. J. Forrester, “The spectrum edge of random matrix ensembles”, *Nucl. Phys. B* **402** (1993), 709–728.
- [12] P. J. Forrester, “Painlevé transcendent evaluation of the scaled distribution of the smallest eigenvalue in the Laguerre orthogonal and symplectic ensembles”, preprint, nlin.SI/0005064, 2000.
- [13] P. J. Forrester and T. D. Hughes, “Complex Wishart matrices and conductance in mesoscopic systems:” exact results. *J. Math. Phys.* **35** (1994), 6736–6747.
- [14] N. R. Goodman, “Statistical analysis based on a certain multivariate complex Gaussian distribution (an introduction)”, *Ann. Math. Stat.* **34** (1963), 152–176.
- [15] J. Gunson, “Proof of a conjecture of Dyson in the statistical theory of energy levels”, *J. Math. Phys.* **4** (1962), 752–753.
- [16] S. Helgason, *Differential geometry and symmetric spaces*, Academic Press, New York, 1962.
- [17] G. Mahoux and M. L. Mehta, “A method of integration over matrix variables IV”, *J. Physique I (France)* **1** (1991), 1093–1108.
- [18] G. Mahoux, M. L. Mehta, and J.-M. Normand, “Matrices coupled in a chain, II: Spacing functions”, *J. Phys. A* **31** (1999), 4457–4464.
- [19] M. L. Mehta, *Random Matrices*, 2nd ed., Academic Press, New York, 1991.
- [20] M. L. Mehta and F. J. Dyson, “Statistical theory of the energy levels of complex systems, V”, *J. Math. Phys.* **4** (1963), 713–719.
- [21] R. J. Muirhead, *Aspects of multivariable statistical theory*. Wiley, New York, 1982.
- [22] T. Nagao and P. J. Forrester, “Asymptotic correlations at the spectrum edge of random matrices”, *Nucl. Phys. B* **435** (1995), 401–420.
- [23] T. Nagao and P. J. Forrester, “The smallest eigenvalue at the spectrum edge of random matrices”, *Nucl. Phys. B* **509** (1998), 561–598.
- [24] T. Nagao and M. Wadati, “Correlation functions of random matrix ensembles related to classical orthogonal polynomials II, III”, *J. Phys. Soc. Japan*, **61** (1992), 78–88, 1910–1918.

- [25] V. Romanovsky, “Sur quelques classes nouvelles de polynômes orthogonaux”, *C. R. Acad. Sci. Paris* **188** (1929), 1023–1025.
- [26] G. Szegő, *Orthogonal polynomials*, 4th ed., Amer. Math. Soc., Providence, RI, 1975.
- [27] C. A. Tracy and H. Widom, “On orthogonal and symplectic matrix ensembles”, *Commun. Math. Phys.* **177** (1996), 727–754.
- [28] C. A. Tracy and H. Widom, “Correlation functions, cluster functions and spacing distributions in random matrices”, *J. Stat. Phys.* **92** (1998), 809–835.
- [29] J. Wishart, “The generalized product moment distribution in samples from a normal multivariate population”, *Biometrika* **20A** (1928), 32–43.
- [30] M. R. Zirnbauer, “Riemannian symmetric superspaces and their origin”, *J. Math. Phys.* **10** (1997), 4986–5018.

PETER J. FORRESTER
DEPARTMENT OF MATHEMATICS AND STATISTICS
UNIVERSITY OF MELBOURNE
PARKVILLE, VICTORIA 3052
AUSTRALIA
p.forrester@ms.unimelb.edu.au

ERIC M. RAINS
AT&T RESEARCH
FLORHAM PARK, NJ 07932
UNITED STATES
rains@research.att.com