

The geography of irregular surfaces

MARGARIDA MENDES LOPES AND RITA PARDINI

We give an overview of irregular complex surfaces of general type, discussing in particular the distribution of the numerical invariants K^2 and χ for minimal ones.

1. Introduction	349
2. Irregular surfaces of general type	350
3. The Castelnuovo–de Franchis inequality	363
4. The slope inequality	364
5. The Severi inequality	368
References	375

1. Introduction

Let S be a minimal surface of general type and let K^2 , χ be its main numerical invariants (Section 2.3). For every pair of positive integers a , b the surfaces with $K^2 = a$, $\chi = b$ belong to finitely many irreducible families, so that in principle their classification is possible. In practice, the much weaker *geographical problem*, i.e., determining the pairs a , b for which there exists a minimal surface of general type with $K^2 = a$ and $\chi = b$, is quite hard.

In the past, the main focus in the study of both the geographical problem and the fine classification of surfaces of general type has been on *regular* surfaces, namely surfaces that have no global 1-forms, or, equivalently, whose first Betti number is 0. The reason for this is twofold: on the one hand, the canonical map of regular surfaces is easier to understand, on the other hand complex surfaces are the main source of examples of differentiable 4-manifolds, hence the simply connected ones are considered especially interesting from that point of view.

So, while, for instance, the geographical problem is by now almost settled and the fine classification of some families of regular surfaces is accomplished

The first author is a member of the Center for Mathematical Analysis, Geometry and Dynamical Systems, IST/UTL and the second author is a member of G.N.S.A.G.A.-I.N.d.A.M. This research was partially supported by the Fundação para a Ciência e Tecnologia (FCT Portugal).

This is an expanded version of the talk given by the second author at the workshop “Classical Algebraic Geometry Today”, MSRI, January 26–30, 2009.

MSC2000: 14J29.

[Barth et al. 1984, §9–11, Chapter VII], little is known about irregular surfaces of general type. In recent years, however, the use of new methods and the revisiting of old ones have produced several new results.

Here we give an overview of these results, with special emphasis on the geographical problem. In addition, we give several examples and discuss some open questions and possible generalizations to higher dimensions (e.g., Theorem 5.2.2).

Notation and conventions. We work over the complex numbers. All varieties are projective algebraic and, unless otherwise specified, smooth. We denote by $J(C)$ the Jacobian of a curve C .

Given varieties X_i and sheaves \mathcal{F}_i on X_i , $i = 1, 2$, we denote by $\mathcal{F}_1 \boxtimes \mathcal{F}_2$ the sheaf $p_1^* \mathcal{F}_1 \otimes p_2^* \mathcal{F}_2$ on $X_1 \times X_2$, where $p_i : X_1 \times X_2 \rightarrow X_i$ is the projection.

2. Irregular surfaces of general type

Unless otherwise specified, a surface is a smooth projective complex surface.

A surface S is of *general type* if the canonical divisor K_S is big. Every surface of general type has a birational morphism onto a unique minimal model, which is characterized by the fact that K_S is nef. A surface (or more generally a variety) S is called *irregular* if the *irregularity* $q(S) := h^0(\Omega_S^1) = h^1(\mathcal{O}_S)$ is > 0 .

2.1. Irrational pencils. A *pencil* on a surface S is a morphism with connected fibers $f : S \rightarrow B$, where B is a smooth curve. A map $\psi : S \rightarrow X$, X a variety, is *composed with a pencil* if there exists a pencil $f : S \rightarrow B$ and a map $\bar{\psi} : B \rightarrow X$ such that $\psi = \bar{\psi} \circ f$. The *genus* of the pencil f is by definition the genus b of B . The pencil f is *irrational* if $b > 0$. Since pullback of forms induces an injective map $H^0(\Omega_B^1) \rightarrow H^0(\Omega_S^1)$, a surface with an irrational pencil is irregular. Clearly, the converse is not true (Remark 2.2.1).

In addition, if the pencil f has genus ≥ 2 , by pulling back two independent 1-forms of B one gets independent 1-forms α and β on S such that $\alpha \wedge \beta = 0$. The following classical result (see [Beauville 1996] for a proof) states that this condition is equivalent to the existence of an irrational pencil of genus ≥ 2 :

Theorem 2.1.1 (Castelnuovo and de Franchis). *Let $\alpha, \beta \in H^0(\Omega_S^1)$ be linearly independent forms such that $\alpha \wedge \beta = 0$. Then there exists a pencil $f : S \rightarrow B$ of genus ≥ 2 and $\alpha_0, \beta_0 \in H^0(\omega_B)$ such that $\alpha = f^* \alpha_0$, $\beta = f^* \beta_0$.*

Let $\alpha_1, \dots, \alpha_k, \beta \in H^0(\Omega_S^1)$ be linearly independent forms such that $\alpha_1, \dots, \alpha_k$ are pullbacks from a curve via a pencil $f : S \rightarrow B$ and $\beta \wedge \alpha_j = 0$ (notice that it is the same to require this for one index j or for all $j = 1, \dots, k$). By Theorem 2.1.1, there exists a pencil $h : S \rightarrow D$ such that, say, α_1 and β are pullbacks of independent 1-forms of D . Let $\psi := f \times h : S \rightarrow B \times D$. Then the forms $\alpha_j \wedge \beta$ are pullbacks of nonzero 2-forms of $B \times D$. Since $\alpha_j \wedge \beta = 0$, it

follows that the image of ψ is a curve C and that $\alpha_1, \dots, \alpha_k, \beta$ are pullbacks of 1-forms of C . This shows that for every $b \geq 2$ there is one-to-one correspondence between pencils of S of genus b and subspaces $W \subset H^0(\Omega_S^1)$ of dimension b such that $\bigwedge^2 W = 0$ and W is maximal with this property.

The existence of a subspace W as above can be interpreted in terms of the cohomology of S with complex coefficients, thus showing that the existence of a pencil of genus ≥ 2 is a topological property [Catanese 1991, Theorem 1.10]. On the contrary, the existence of pencils of genus 1 is not detected by the topology (Remark 2.2.1).

A classical result of Severi (see [Severi 1932; Samuel 1966]) states that a surface of general type has finitely many pencils of genus ≥ 2 . However a surface of general type can have infinitely many pencils of genus 1, as it is shown by the following example:

Example 2.1.2. Let E be a curve of genus 1, $O \in E$ a point, $A := E \times E$ and $L := \mathbb{C}_A(\{O\} \times E + E \times \{O\})$. For every $n \geq 1$, the map $h_n : A \rightarrow E$ defined by $(x, y) \mapsto x + ny$ is a pencil of genus 1. Let now $S \rightarrow A$ be a double cover branched on a smooth ample curve $D \in |2dL|$ (see Section 2.4(d) for a quick review of double covers). The surface S is minimal of general type, and for every n the map h_n induces a pencil $f_n : S \rightarrow E$. The general fiber of f_n is a double cover of an elliptic curve isomorphic to E , branched on $2d(n^2 + 1)$ points, hence it has genus $dn^2 + d + 1$. It follows that the pencils f_n are all distinct.

2.2. The Albanese map. The *Albanese variety* of S is defined as $\text{Alb}(S) := H^0(\Omega_S^1)^\vee / H_1(S, \mathbb{Z})$. By Hodge theory, $\text{Alb}(S)$ is a compact complex torus and, in addition, it can be embedded in projective space, namely it is an abelian variety. For a fixed base point $x_0 \in S$ one defines the Albanese morphism $a_{x_0} : S \rightarrow \text{Alb}(S)$ by $x \mapsto \int_{x_0}^x -$; see [Beauville 1996, Chapter V]. Choosing a different base point in S , the Albanese morphism just changes by a translation of $\text{Alb}(S)$, so we often ignore the base point and just write a . By construction, the map $a_* : H_1(S, \mathbb{Z}) \rightarrow H_1(\text{Alb}(S), \mathbb{Z})$ is surjective, with kernel equal to the torsion subgroup of $H_1(S, \mathbb{Z})$, and the map $a^* : H^0(\Omega_{\text{Alb}(S)}^1) \rightarrow H^0(\Omega_S^1)$ is an isomorphism, so if $q(S) > 0$ it follows immediately that a is nonconstant. The dimension of $a(S)$ is called the *Albanese dimension* of S and it is denoted by $\text{Albdim}(S)$. S is of *Albanese general type* if $\text{Albdim}(S) = 2$ and $q(S) > 2$.

The morphism $a : S \rightarrow \text{Alb}(S)$ is characterized up to unique isomorphism by the following universal property: for every morphism $S \rightarrow T$, with T a complex torus, there exists a unique factorization $S \xrightarrow{a} \text{Alb}(S) \rightarrow T$. It follows immediately that the image of a generates $\text{Alb}(S)$, namely that $a(S)$ is not contained in any proper subtorus of $\text{Alb}(S)$. Using the Stein factorization and the fact that for a smooth curve B the Abel–Jacobi map $B \rightarrow J(B)$ is an embedding, one can

show that if $\text{Albdim}(S) = 1$, then $B := a(S)$ is a smooth curve of genus $q(S)$ and the map $a : S \rightarrow B$ has connected fibers. In this case, the map $a : S \rightarrow B$ is called the *Albanese pencil* of S . By the analysis of Section 2.1, $\text{Albdim}(S) = 1$ (“ a is composed with a pencil”) if and only if $\alpha \wedge \beta = 0$ for every pair of 1-forms $\alpha, \beta \in H^0(\Omega_S^1)$ if and only if there exists a surjective map $f : S \rightarrow B$ where B is a smooth curve of genus $q(S)$. In this case, if $q(S) > 1$ then f coincides with the Albanese pencil.

Since, as we recalled in Section 2.1, the existence of a pencil of given genus ≥ 2 is a topological property of S , the Albanese dimension of a surface is a topological property.

Remark 2.2.1. Let $f : S \rightarrow B$ be an irrational pencil. By the universal property of the Albanese variety, there is a morphism of tori $\text{Alb}(S) \rightarrow J(B)$. The differential of this morphism at 0 is dual to $f^* : H^0(\omega_B) \rightarrow H^0(\Omega_S^1)$, hence it is surjective and $\text{Alb}(S) \rightarrow J(B)$ is surjective, too. So if $\text{Albdim}(S) = 2$ and $\text{Alb}(S)$ is simple, S has no irrational pencil. It is easy to produce examples of this situation, for instance by considering surfaces that are complete intersections inside a simple abelian variety; see Section 2.4(c).

If $\text{Albdim}(S) = q(S) = 2$, S has an irrational pencil if and only if there exists a surjective map $\text{Alb}(S) \rightarrow E$, where E is an elliptic curve. So, by taking double covers of principally polarized abelian surfaces branched on a smooth ample curve — see Section 2.4(d) — one can construct a family of minimal surfaces of general type such that the general surface in the family has no irrational pencil but some special ones have.

If $\text{Albdim}(S) = 2$, then a contracts finitely many irreducible curves. By Grauert’s criterion ([Barth et al. 1984, Theorem III.2.1]), the intersection matrix of the set of curves contracted by a is negative definite. An irreducible curve C of S is contracted by a if and only if the restriction map $H^0(\Omega_S^1) \rightarrow H^0(\omega_{C^\nu})$ is trivial, where $C^\nu \rightarrow C$ is the normalization. So, every rational curve of S is contracted by a . More generally, by the universal property the map $C^\nu \rightarrow \text{Alb}(S)$ factorizes through $J(C^\nu) = \text{Alb}(C^\nu)$, hence $a(C)$ spans an abelian subvariety of $\text{Alb}(S)$ of dimension at most $g(C^\nu)$. So, for instance, if $\text{Alb}(S)$ is simple every curve of S of geometric genus $< q(S)$ is contracted by a .

An irreducible curve C of S such that $a(C)$ spans a proper abelian subvariety $T \subset \text{Alb}(S)$ has $C^2 \leq 0$. In particular, if C has geometric genus $< q(S)$ then $C^2 \leq 0$. Indeed, consider the nonconstant map $\bar{a} : S \rightarrow A/T$ induced by a . If the image of \bar{a} is a surface, then C is contracted to a point, hence $C^2 < 0$. If the image of \bar{a} is a curve, then C is contained in a fiber, hence by Zariski’s lemma ([Barth et al. 1984, Lem.III.8.9]) $C^2 \leq 0$ and $C^2 = 0$ if and only if C moves in an irrational pencil. In view of the fact that S has finitely many irrational pencils

of genus ≥ 2 , the irreducible curves of S whose image via a spans an abelian subvariety of $\text{Alb}(S)$ of codimension > 1 belong to finitely many numerical equivalence classes.

We close this section by giving an example that shows that the degree of the Albanese map of a surface with $\text{Albdim} = 2$ is not a topological invariant.

Example 2.2.2. We describe an irreducible family of smooth minimal surfaces of general type such that the Albanese map of the general element of the family is generically injective but for some special elements the Albanese map has degree 2 onto its image. The examples are constructed as divisors in a double cover $p : V \rightarrow A$ of an abelian threefold A .

Let A be an abelian threefold and let L be an ample line bundle of A such that $|2L|$ contains a smooth divisor D . There is a double cover $p : V \rightarrow A$ branched on D and such that $p_*\mathcal{O}_V = \mathcal{O}_A \oplus L^{-1}$; see Section 2.4(d). The variety V is smooth and, arguing as in Section 2.4(d), one shows that the Albanese map of V coincides with p . Notice that this implies that the map $p_* : H_1(V, \mathbb{Z}) \rightarrow H_1(A, \mathbb{Z})$ is an isomorphism up to torsion (see Section 2.2).

Let now $Y \subset A$ be a very ample divisor such that $h^0(\mathcal{O}_A(Y) \otimes L^{-1}) > 0$ and set $X := p^*Y$. If Y is general, then both X and Y are smooth. Let now $X' \in |X|$ be a smooth element. By the adjunction formula, X' is smooth of general type. By the Lefschetz theorem on hyperplane sections, the inclusion $X' \rightarrow V$ induces an isomorphism $\pi_1(X') \simeq \pi_1(V)$, which in turn gives an isomorphism $H_1(X', \mathbb{Z}) \simeq H_1(V, \mathbb{Z})$. Composing with $p_* : H_1(V, \mathbb{Z}) \rightarrow H_1(A, \mathbb{Z})$ we get an isomorphism (up to torsion) $H_1(X', \mathbb{Z}) \rightarrow H_1(A, \mathbb{Z})$, which is induced by the map $p|_{X'} : X' \rightarrow A$. Hence $p|_{X'} : X' \rightarrow A$ is the Albanese map of X' . Now the map $p|_{X'}$ has degree 2 onto its image if X' is invariant under the involution σ associated to p , and it is generically injective otherwise. By the projection formula for double covers, the general element of $|X|$ is not invariant under σ if and only if $h^0(\mathcal{O}_A(Y) \otimes L^{-1}) > 0$, hence we have the required example.

2.3. Numerical invariants and geography. To a minimal complex surface S of general type, one can attach several integer invariants, besides the irregularity $q(S) = h^0(\Omega_S^1) > 0$:

- the self intersection K_S^2 of the canonical class,
- the *geometric genus* $p_g(S) := h^0(K_S) = h^2(\mathcal{O}_S)$,
- the *holomorphic Euler–Poincaré characteristic*, $\chi(S) := h^0(\mathcal{O}_S) - h^1(\mathcal{O}_S) + h^2(\mathcal{O}_S) = 1 - q(S) + p_g(S)$,
- the second Chern class $c_2(S)$ of the tangent bundle, which coincides with the topological Euler characteristic of S .

All these invariants are determined by the topology of S plus the orientation induced by the complex structure. Indeed (see [Barth et al. 1984, I.1.5]), by Noether's formula we have:

$$K_S^2 + c_2(S) = 12\chi(S), \quad (2.3.1)$$

and by the Thom–Hirzebruch index theorem:

$$\tau(S) = 2(K_S^2 - 8\chi(S)), \quad (2.3.2)$$

where $\tau(S)$ denotes the index of the intersection form on $H^2(S, \mathbb{C})$, namely the difference between the number of positive and negative eigenvalues. So K_S^2 and $\chi(S)$ are determined by the (oriented) topological invariants $c_2(S)$ and $\tau(S)$. By Hodge theory the irregularity $q(S)$ is equal to $\frac{1}{2}b_1(S)$, where $b_1(S)$ is the first Betti number. So $q(S)$ is also determined by the topology of S and the same is true for $p_g(S) = \chi(S) + q(S) - 1$.

It is apparent from the definition that these invariants are not independent. So it is usual to take $K_S^2, \chi(S)$ (or, equivalently, $K_S^2, c_2(S)$) as the main numerical invariants. These determine the Hilbert polynomial of the n -canonical image of S for $n \geq 2$, and by a classical result [Gieseker 1977] the coarse moduli space $\mathcal{M}_{a,b}$ of surfaces of general type with $K^2 = a, \chi = b$ is a quasiprojective variety. Roughly speaking, this means that surfaces with fixed K^2 and χ are parametrized by a finite number of irreducible varieties, hence in principle they can be classified. In practice, however, the much more basic *geographical question*, i.e., “for what values of a, b is $\mathcal{M}_{a,b}$ nonempty?” is already nontrivial.

The invariants K^2, χ are subject to the following restrictions:

- $K^2, \chi > 0$,
- $K^2 \geq 2\chi - 6$ (Noether's inequality),
- $K^2 \leq 9\chi$ (Bogomolov–Miyaoka–Yau inequality).

All these inequalities are sharp and it is known that for “almost all” a, b in the admissible range the space $\mathcal{M}_{a,b}$ is nonempty. (The possible exceptions seem to be due to the method of proof and not to the existence of special areas in the admissible region for the invariants of surfaces of general type).

In this note we focus on the geographical question for irregular surfaces. More precisely, we address the following questions:

“for what values of a, b does there exist a minimal surface of general type S with $K^2 = a, \chi = b$ such that:

- $q(S) > 0$?”
- S has an irrational pencil?”
- $\text{Albdim}(S) = 2$?”
- $q(S) > 0$ and S has no irrational pencil?”

Remark 2.3.1. If $S' \rightarrow S$ is an étale cover of degree d and S is minimal of general type, S' is also minimal of general type with invariants $K_{S'}^2 = dK_S^2$ and $\chi(S') = d\chi(S)$. The first three properties listed above are stable under étale covers. Since the first Betti number of a surface is equal to $2q$, an irregular surface has étale covers of degree d for any $d > 0$. Hence, if for some a, b the answer to one of these three questions is affirmative, the same is true for all the pairs da, db , with d a positive integer.

The main known inequalities for the invariants of irregular surfaces of general type are illustrated in the following sections. Here we only point out the following simple consequence of Noether’s inequality:

Proposition 2.3.2. *Let S be a minimal irregular surface of general type. Then:*

$$K_S^2 \geq 2\chi(S).$$

Proof. Assume for contradiction that $K_S^2 < 2\chi(S)$. Then an étale cover S' of degree $d \geq 7$ has $K_{S'}^2 < 2\chi(S') - 6$, violating Noether’s inequality. \square

More generally, the following inequality holds for minimal irregular surfaces of general type [Debarre 1982]:

$$K^2 \geq \max\{2p_g, 2p_g + 2(q - 4)\}. \tag{2.3.3}$$

The inequality (2.3.3) implies that irregular surfaces with $K^2 = 2\chi$ have $q = 1$. These surfaces are described in Section 2.5(b).

2.4. Basic constructions. Some constructions of irregular surfaces of general type have already been presented in the previous sections. We list and describe briefly the most standard ones:

(a) Products of curves. Take $S := C_1 \times C_2$, with C_i a curve of genus $g_i \geq 2$. S has invariants

$$K^2 = 8(g_1 - 1)(g_2 - 1), \quad \chi = (g_1 - 1)(g_2 - 1), \quad q = g_1 + g_2, \quad p_g = g_1g_2.$$

In particular these surfaces satisfy $K^2 = 8\chi$. The Albanese variety is the product $J(C_1) \times J(C_2)$ and the Albanese map induces an isomorphism onto its image. The two projections $S \rightarrow C_i$ are pencils of genus $g_i \geq 2$.

(b) Symmetric products. Take $S := S^2C$, where C is a smooth curve of genus $g \geq 3$. Consider the natural map $p : C \times C \rightarrow S$, which is the quotient map by the involution ι that exchanges the two factors of $C \times C$. The ramification divisor of p is the diagonal $\Delta \subset C \times C$, hence we have:

$$K_{C \times C} = p^*K_S + \Delta.$$

Computing intersections on $C \times C$ we get

$$K_S^2 = (g - 1)(4g - 9).$$

Global 1 and 2-forms on S^2C correspond to forms on $C \times C$ that are invariant under ι . Writing down the action of ι on $H^0(\Omega_{C \times C}^i)$, one obtains canonical identifications:

$$H^0(\omega_S) = \wedge^2 H^0(\omega_C), \quad H^0(\Omega_S^1) = H^0(\omega_C). \tag{2.4.1}$$

Thus we have:

$$p_g = g(g - 1)/2, \quad q = g, \quad \chi = g(g - 3)/2 + 1.$$

Since $p_g(S) > 0$ and $K_S^2 > 0$, it follows that S is of general type. Notice that by Theorem 2.1.1 S has no irrational pencil of genus ≥ 2 , since by (2.4.1) the natural map $\wedge^2 H^0(\Omega_S^1) \rightarrow H^0(\omega_S)$ is injective.

The points of S can be identified with the effective divisors of degree 2 of C . If C is hyperelliptic, then the g_2^1 of C gives a smooth rational curve Γ of S such that $\Gamma^2 = 1 - g$. Let $(P_0, Q_0) \in C \times C$ be a point: the map $C \times C \rightarrow J(C) \times J(C)$ defined by $(P, Q) \mapsto (P - P_0, Q - Q_0)$ is the Albanese map of $C \times C$ with base point (P_0, Q_0) . Composing with the addition map, one obtains a map $C \times C \rightarrow J(C)$ that is invariant for the action of ι and therefore induces a map $a : S \rightarrow J(C)$, which can be written explicitly as $P + Q \mapsto P + Q - P_0 - Q_0$. Using the universal property, one shows that a is the Albanese map of S . By the Riemann–Roch theorem, if C is not hyperelliptic a is injective, while if C is hyperelliptic a contracts Γ to a point and is injective on $S \setminus \Gamma$. Since $H^0(\omega_S)$ is the pullback of $H^0(\Omega_{J(C)}^2) = \wedge^2 H^0(\omega_C)$ via the Albanese map a , the points of S where the differential of a fails to be injective are precisely the base points of $|K_S|$. So, if C is not hyperelliptic then a is an isomorphism of S with its image and if C is hyperelliptic, then a gives an isomorphism of $S \setminus \Gamma$ with its image.

Notice that as g goes to infinity, the ratio $K_S^2/\chi(S)$ approaches 8 from below.

(c) Complete intersections. Let V be an irregular variety of dimension $k+2 \geq 3$. For instance, one can take as V an abelian variety or a product of curves not all rational. Given $|D_1|, \dots, |D_k|$ free and ample linear systems on V such that $K_V + D_1 + \dots + D_k$ is nef and big, we take

$$S = D_1 \cap \dots \cap D_k,$$

with $D_i \in |D_i|$ general, so that S is smooth. By the adjunction formula, K_S is the restriction to S of $K_V + D_1 + \dots + D_k$, hence S is minimal of general type. Since the D_i are ample, the Lefschetz Theorem for hyperplane sections gives an isomorphism $H_1(S, \mathbb{Z}) \cong H_1(V, \mathbb{Z})$. Hence the Albanese map of S is just the restriction of the Albanese map of V .

The numerical invariants of S can be computed by means of standard exact sequences on V . If $k = 1$ and $D_1 \in |rH|$, where H is a fixed ample divisor, one has:

$$K_S^2 = r^3 H^3 + O(r^2), \quad \chi(S) = r^3 H^3 / 6 + O(r^2),$$

so the ratio $K_S^2 / \chi(S)$ tends to 6 as r goes to infinity. Similarly, for $k = 2$ and $D_1, D_2 \in |rH|$, one has:

$$K_S^2 = 4r^4 H^4 + O(r^3), \quad \chi(S) = 7r^4 H^4 / 12 + O(r^3),$$

and $K_S^2 / \chi(S)$ tends to $48/7$ as r goes to infinity.

(d) Double covers. If Y is an irregular surface, any surface S that dominates Y is irregular, too, and $\text{Albdim}(S) \geq \text{Albdim}(Y)$. The simplest instance of this situation in which the map $S \rightarrow Y$ is not birational is that of a double cover. A smooth double cover of a variety Y is determined uniquely by a line bundle L on Y and a smooth divisor $D \in |2L|$. Set $\mathcal{E} := \mathcal{O}_Y \oplus L^{-1}$ and let \mathbb{Z}_2 act on \mathcal{E} as multiplication by 1 on \mathcal{O}_Y and multiplication by -1 on L^{-1} . To define on \mathcal{E} an \mathcal{O}_Y -algebra structure compatible with this \mathbb{Z}_2 -action it suffices to give a map $\mu : L^{-2} \rightarrow \mathcal{O}_Y$: we take μ to be a section whose zero locus is D , set $S := \text{Spec } \mathcal{E}$ and let $\pi : S \rightarrow Y$ be the natural map. S is easily seen to be smooth if and only if D is. By construction, one has:

$$H^i(\mathcal{O}_S) = H^i(\mathcal{O}_Y) \oplus H^i(L^{-1}).$$

In particular, if L is nef and big then by Kawamata–Viehweg vanishing $H^1(\mathcal{O}_S) = H^1(\mathcal{O}_Y)$, hence the induced map $\text{Alb}(S) \rightarrow \text{Alb}(Y)$ is an isogeny. It is actually an isomorphism: since $H^0(\Omega_S^1) = H^0(\Omega_Y^1)$, the induced \mathbb{Z}_2 action on $\text{Alb}(S)$ is trivial. Since $D = 2L$ is nef and big and effective, it is nonempty and therefore we may choose a base point $x_0 \in S$ that is fixed by \mathbb{Z}_2 . The Albanese map with base point x_0 is \mathbb{Z}_2 -equivariant, hence it descends to a map $Y \rightarrow \text{Alb}(S)$. So by the universal property there is a morphism $\text{Alb}(Y) \rightarrow \text{Alb}(S)$ which is the inverse of the morphism $\text{Alb}(S) \rightarrow \text{Alb}(Y)$ induced by π .

A local computation gives the following pullback formula for the canonical divisor:

$$K_S = \pi^*(K_Y + L).$$

By this formula, if $K_Y + L$ is nef and big the surface S is minimal of general type. The numerical invariants of S are:

$$K_S^2 = 2(K_Y + L)^2, \quad \chi(S) = 2\chi(Y) + L(K_Y + L)/2.$$

Hence for “large” L , the ratio $K_S^2 / \chi(S)$ tends to 4.

If Y is an abelian surface and L is ample, one has $\text{Albdim}(S) = 2$ and $K_S^2 = 4\chi(S)$.

2.5. Examples. The constructions of irregular surfaces of Section 2.4 can be combined to produce more sophisticated examples; see Example 2.2.2, for instance. However the computations of the numerical invariants suggest that in infinite families of examples the ratio K^2/χ converges, so that it does not seem easy to fill by these methods large areas of the admissible region for the invariants K^2, χ .

Here we collect some existence results for irregular surfaces.

(a) $\chi = 1$. As explained in Section 2.3, $\chi = 1$ is the smallest possible value for a surface of general type. Since $K^2 \leq 9\chi$ (Section 2.3), in this case we have $K^2 \leq 9$, hence by (2.3.3) $q = p_g \leq 4$. To our knowledge, the only known example of an irregular surface of general type with $K^2 = 9, \chi = 1$ was recently constructed by Donald Cartwright and Tim Steiger (unpublished). It has $q = 1$.

We recall briefly what is known about the classification of these surfaces for the possible values of q .

$q = 4$: S is the product of two curves of genus 2 by Theorem 3.0.4, due to Beauville. So $K^2 = 8$ in this case.

$q = 3$: By [Hacon and Pardini 2002] and [Pirola 2002] (see also [Catanese et al. 1998]) these surfaces belong to two families. They are either the symmetric product S^2C of a curve C of genus 3 ($K^2 = 6$) or free \mathbb{Z}_2 -quotients of a product of curves $C_1 \times C_2$ where $g(C_1) = 2, g(C_2) = 3$ ($K^2 = 8$).

$q = 2$: Surfaces with $p_g = q = 2$ having an irrational pencil (hence in particular those with $\text{Albdim}(S) = 1$) are classified in [Zucconi 2003]. They have either $K^2 = 4$ or $K^2 = 8$.

Let (A, Θ) be a principally polarized abelian surface A . A double cover $S \rightarrow A$ branched on a smooth curve of $|2\Theta|$ is a minimal surface of general type with $K^2 = 4, p_g = q = 2$ and it has no irrational pencil if and only if A is simple; see Section 2.4(d). In [Ciliberto and Mendes Lopes 2002] it is proven that this is the only surface with $p_g = q = 2$ and nonbirational bicanonical map that has no pencil of curves of genus 2. An example with $p_g = q = 2, K^2 = 5$ and no irrational pencil is constructed in [Chen and Hacon 2006].

$q = 1$: For S a minimal surface of general type with $p_g = q = 1$, we denote by E the Albanese curve of S and by g the genus of the general fiber of the Albanese pencil $a : S \rightarrow E$.

The case $K^2 = 2$ is classified in [Catanese 1981]. These surfaces are constructed as follows. Let E be an elliptic curve with origin O . The map $E \times E \rightarrow E$ defined by $(P, Q) \mapsto P + Q$ descends to a map $S^2E \rightarrow E$ whose fibers are smooth rational curves. We denote by F the algebraic equivalence class of a fiber of $S^2E \rightarrow E$. The curves $\{P\} \times E$ and $E \times \{P\}$ map to curves

$D_P \subset S^2E$ such that $D_P F = D_P^2 = 1$. The curves D_P , $P \in A$, are algebraically equivalent and $h^0(D_P) = 1$. We denote by D the algebraic equivalence class of D_P . Clearly D and F generate the Néron–Severi group of S^2E . All the surfaces S are (minimal desingularizations of) double covers of S^2E branched on a divisor B numerically equivalent to $6D - 2F$ and with at most simple singularities. The composite map $S \rightarrow S^2E \rightarrow E$ is the Albanese pencil of X and its general fiber has genus 2.

The case $K^2 = 3$ is studied in [Catanese and Ciliberto 1991; 1993]. One has either $g = 2$ or $g = 3$. If $g = 2$, then S is birationally a double cover of S^2E , while if $g = 3$ S is birational to a divisor in S^3E . For $K^2 = 4$, several components of the moduli space are constructed in [Pignatelli 2009] (all these examples have $g = 2$). Rito [2007; 2010b; 2010a] gave examples with $K^2 = 2, \dots, 8$. The case in which S is birational to a quotient $(C \times F)/G$, where C and F are curves and G is a finite group is considered in [Carnovale and Polizzi 2009; Mistretta and Polizzi 2010; Polizzi 2008; 2009]: when $(C \times F)/G$ has at most canonical singularities the surface S has $K^2 = 8$, but there are also examples with $K^2 = 2, 3, 5$.

(b) The line $K^2 = 2\chi$. As pointed out in Proposition 2.3.2, for irregular surfaces the lower bound for the ratio K^2/χ is 2. Irregular surfaces attaining this lower bound were studied in [Horikawa 1977; 1981]. Their structure is fairly simple: they have $q = 1$, the fibers of the Albanese pencil $a : S \rightarrow E$ have genus 2 (compare Proposition 4.1.4) and the quotient of the canonical model of S by the hyperelliptic involution is a \mathbb{P}^1 -bundle over E . The moduli space of these surfaces is studied in [Horikawa 1981].

We just show here that for every integer $d > 0$ there exists a minimal irregular surface of general type with $K^2 = 2\chi$ and $\chi = d$. In (a) above we have sketched the construction of such a surface S with $K_S^2 = 2$, $\chi(S) = 1$. Let $a : S \rightarrow E$ be the Albanese pencil and let $E' \rightarrow E$ be an unramified cover of degree d . Then the map $S' \rightarrow S$ obtained from $E' \rightarrow E$ by taking base change with $S \rightarrow E$ is a connected étale cover and S' is minimal of general type with $K^2 = 2d$, $\chi = d$. By construction S' maps onto E' , hence $q(S') > 0$. By Proposition 4.1.4 we have $q(S') = 1$, hence $S' \rightarrow E'$, having connected fibers, is the Albanese pencil of S' .

Alternatively, here is a direct construction for χ even. Let $Y = \mathbb{P}^1 \times E$, with E an elliptic curve and let $L := \mathcal{O}_{\mathbb{P}^1}(3) \boxtimes \mathcal{O}_E(kO)$, where $k \geq 1$ is an integer and $O \in E$ is a point. Let $D \in |2L|$ be a smooth curve and let $\pi : S \rightarrow Y$ be the double cover given by the relation $2L \equiv D$; see Section 2.4(d). The surface is smooth, since D is smooth, and it is minimal of general type since $K_S = \pi^*(K_Y + L) = \pi^*(\mathcal{O}_{\mathbb{P}^1}(1) \boxtimes \mathcal{O}_E(kO))$ is ample. The invariants are

$$K^2 = 4k, \chi = 2k.$$

One shows as above that $q(S) = 1$ and $S \rightarrow E$ is the Albanese pencil.

(c) Surfaces with an irrational pencil with general fiber of genus g . We use the same construction as in the previous case. Let $g \geq 2$. Take $Y = \mathbb{P}^1 \times E$ with E an elliptic curve, Δ a divisor of positive degree of E and set $L := \mathcal{O}_{\mathbb{P}^1}(g + 1) \boxtimes \mathcal{O}_E(\Delta)$. Let $D \in |2L|$ be a smooth curve and $\pi : S \rightarrow Y$ the double cover given by the relation $2L \equiv D$. S is smooth minimal of general type, with invariants

$$K_S^2 = 4(g - 1) \deg \Delta, \quad \chi(S) = g \deg \Delta.$$

The projection $Y \rightarrow E$ lifts to a pencil $S \rightarrow E$ of hyperelliptic curves of genus g . Here $K_S^2/\chi(S)$ is equal to $4(g - 1)/g$, which is the lowest possible value by Theorem 4.1.3.

(d) The line $K^2 = 9\chi$. By [Miyaoka 1984, Theorem 2.1] minimal surfaces of general type with $K^2 = 9\chi$ have ample canonical class. By Yau’s results [1977], surfaces with $K^2 = 9\chi$ and ample canonical class are quotients of the unit ball in \mathbb{C}^2 by a discrete subgroup. The existence of several examples has been shown using this description [Barth et al. 1984, §9, Chapter VII].

Three examples have been constructed in [Hirzebruch 1983] as Galois covers of the plane branched on an arrangement of lines.

For later use, we sketch here one of these constructions. Let $P_1, \dots, P_4 \in \mathbb{P}^2$ be points in general positions and let L_1, \dots, L_6 be equations for the lines through P_1, \dots, P_4 . Let $X \rightarrow \mathbb{P}^2$ be the normal finite cover corresponding to the field inclusion

$$\mathbb{C}(\mathbb{P}^2) \subset \mathbb{C}(\mathbb{P}^2)((L_1/L_6)^{\frac{1}{5}}, \dots, (L_5/L_6)^{\frac{1}{5}}).$$

The cover $X \rightarrow \mathbb{P}^2$ is abelian with Galois group \mathbb{Z}_5^5 and one can show, for instance by the methods of [Pardini 1991], that X is singular over P_1, \dots, P_4 and that the cover $S \rightarrow \hat{\mathbb{P}}^2$ obtained by blowing up P_1, \dots, P_4 and taking base change and normalization is smooth. The cover $S \rightarrow \hat{\mathbb{P}}^2$ is branched of order 5 on the union B of the exceptional curves and of the strict transforms of the lines L_j . Hence the canonical class K_S is numerically equivalent to the pull back of $\frac{1}{5}(9L - 3(E_1 + \dots + E_4))$, where L is the pullback on $\hat{\mathbb{P}}^2$ class of a line in \mathbb{P}^2 and the E_i are the exceptional curves of $\hat{\mathbb{P}}^2 \rightarrow \mathbb{P}^2$. It follows that K_S is ample and $K_S^2 = 3^2 \cdot 5^4$. The divisor B has 15 singular points, that are precisely the points of $\hat{\mathbb{P}}^2$ whose preimage consists of 5^3 points. Hence, denoting by e the topological Euler characteristic of a variety, we have

$$c_2(S) = e(S) = 5^5[e(\hat{\mathbb{P}}^2) - e(B)] + 5^4[e(B) - 15] + 5^3 \cdot 15 = 3 \cdot 5^4.$$

Thus S satisfies $K^2 = 3c_2$ or, equivalently, $K^2 = 9\chi$. In [Ishida 1983] it is shown that the irregularity $q(S)$ is equal to 30. To prove that $\text{Albdim } S = 2$, by the discussion in Section 2.1 it is enough to show that S has more than one irrational pencil. The surface $\hat{\mathbb{P}}^2$ has 5 pencils of smooth rational curves, induced by the systems h_i of lines through each of the P_i , $i = 1, \dots, 4$, and by the system h_5 of conics through P_1, \dots, P_4 . For $i = 1, \dots, 5$, denote by $f_i : S \rightarrow B_i$ the pencil induced by h_i and denote by F_i the general fiber of f_i . For $i = 1, \dots, 5$, the subgroup $H_i < \mathbb{Z}_5^5$ that maps F_i to itself has order 5^3 and the restricted cover $F_i \rightarrow \mathbb{P}^1$ is branched at 4 points. So F_i has genus 76 by the Hurwitz formula. There is a commutative diagram

$$\begin{array}{ccc}
 S & \longrightarrow & \hat{\mathbb{P}}^2 \\
 f_i \downarrow & & \downarrow \\
 B_i & \longrightarrow & \mathbb{P}^1
 \end{array} \tag{2.5.1}$$

where the map $B_i \rightarrow \mathbb{P}^1$ is an abelian cover with Galois group $\mathbb{Z}_5^5/H_i \cong \mathbb{Z}_5^2$. The branch points of $B_i \rightarrow \mathbb{P}^1$ correspond to the multiple fibers of f_i , hence there are 3 of them and B_i has genus 6 by the Hurwitz formula. One computes $F_i F_j = 5$ for $i \neq j$, hence the pencils F_i are all distinct.

Since the group $H_i \cap H_j$ acts faithfully on the set $F_i \cap F_j$ for F_i, F_j general, it follows that $H_i \cap H_j$ has order 5 and $H_i + H_j = \mathbb{Z}_5^5$. We use this remark to show that $H^0(\Omega_S^1) = \bigoplus_{i=1}^5 V_i$, where $V_i := f_i^* H^0(\omega_{B_i})$, and therefore that $\text{Alb}(S)$ is isogenous to $J(B_1) \times \dots \times J(B_5)$. Since $q(S) = 30$ and $\dim V_i = 6$, it is enough to show that the V_i are in direct sum in $H^0(\Omega_S^1)$. Each subspace V_i decomposes under the action of \mathbb{Z}_5^5 as a direct sum of eigenspaces relative to some subset of the group of characters $\text{Hom}(\mathbb{Z}_5^5, \mathbb{C}^*)$. Notice that the trivial character never occurs in the decomposition since $B_i/\mathbb{Z}_5^5 = \mathbb{P}^1$. The diagram (2.5.1) shows that the characters occurring in the decomposition of V_i belong to H_i^\perp . Since $H_i + H_j = \mathbb{Z}_5^5$ for $i \neq j$, it follows that each character occurs in the decompositions of at most one the V_i , and therefore that there is no linear relation among the V_i .

(e) The ratio K^2/χ . Sommese [1984] showed that the ratios $K^2(S)/\chi(S)$, for S a minimal surface of general type, form a dense set in the admissible interval $[2, 9]$. His construction can be used to prove:

Proposition 2.5.1. (i) *The ratios $K_S^2/\chi(S)$, as S ranges among surfaces with $\text{Albdim}(S) = 1$, are dense in the interval $[2, 8]$.*

(ii) *The ratios $K_S^2/\chi(S)$, as S ranges among surfaces with $\text{Albdim}(S) = 2$, are dense in the interval $[4, 9]$.*

Proof. Let X be a minimal surface of general type and let $f : X \rightarrow B$ be an irrational pencil. Denote by $g \geq 2$ the genus of a general fiber of f and write K^2, χ for $K_X^2, \chi(X)$. Given positive integers d, k , we construct a surface $S_{d,k}$ as follows:

(1) We take an unramified degree d cover $B' \rightarrow B$ and let $Y_d \rightarrow X$ be the cover obtained by taking base change with $f : X \rightarrow B$.

(2) We take a double cover $B'' \rightarrow B'$ branched on $2k > 0$ general points and let $S_{d,k} \rightarrow Y_d$ be the cover obtained from $B'' \rightarrow B'$ by base change.

The étale cover $Y_d \rightarrow X$ is connected, hence Y_d is a minimal surface of general type with $K_{Y_d}^2 = dK^2$ and $\chi(Y_d) = d\chi$. By Section 2.4(d), the surface $S_{d,k}$ is smooth, since the branch points of $B'' \rightarrow B'$ are general, and it is minimal of general type since $K_{S_{d,k}}$ is numerically the pullback of $K_{Y_d} + kF$, where F is a fiber of $Y_d \rightarrow B'$ (F is the same as the general fiber of $X \rightarrow B$). By the formulae for double covers we have

$$\frac{K_{S_{d,k}}^2}{\chi(S_{d,k})} = \frac{2dK^2 + 8k(g - 1)}{2d\chi + k(g - 1)} = \frac{K^2}{\chi} + \left(8 - \frac{K^2}{\chi}\right) \frac{k(g - 1)}{2d\chi + k(g - 1)}. \tag{2.5.2}$$

This formula shows that the ratio $K_{S_{d,k}}^2/\chi(S_{d,k})$ is in the interval $[8, K^2/\chi]$ if $K^2 \geq 8\chi$ and it is in $[K^2/\chi, 8]$ otherwise. It is not difficult to show that as d, k vary one obtains a dense set in the appropriate interval [Sommese 1984].

Now to prove the statement it is enough to apply the construction to suitable surfaces. If one takes X to be the surface with $K^2 = 9\chi$ described in (c) and $f : X \rightarrow B$ one of the 5 irrational pencils of X , then the surfaces $S_{d,k}$ have Albanese dimension 2 and the ratios of their numerical invariants are dense in $[8, 9]$.

If one takes X to be a double cover of $E \times E$ branched on a smooth ample curve as in Section 2.4(d) and $f : X \rightarrow E$ one of the induced pencil, then the surfaces $S_{d,k}$ have Albanese dimension 2 and the ratio of their numerical invariants are dense in $[4, 8]$.

Finally, we take X an irregular surface with $K^2 = 2\chi$. Since $q(X) = 1$ (see (c) above), we can take $f : X \rightarrow B$ to be the Albanese pencil. In this case the ratios of the numerical invariants of the surfaces $S_{d,k}$ are dense in the interval $[2, 8]$. To complete the proof we show that the surfaces $S_{d,k}$ have Albanese dimension 1. The surfaces Y_d satisfy $K^2 = 2\chi$, hence they also have $q = 1$. The induced pencil $S_{d,k} \rightarrow B''$ has genus $k + 1$, so we need to show that the irregularity of $S_{d,k}$ is equal to $k + 1$. Denote by L the line bundle of Y_d associated to the double cover $S_{d,k} \rightarrow Y_d$. By construction $L = \mathcal{O}_{Y_d}(F_1 + \dots + F_k)$, where the F_i are fibers of the Albanese pencil, and if $k > 1$ we can take the F_i smooth and distinct. We have $q(S_{d,k}) = q(Y_d) + h^1(L^{-1}) = 1 + h^1(L^{-1})$. Finally, $h^1(L^{-1}) = k$ can be

proven using the restriction sequence

$$0 \rightarrow L^{-1} \rightarrow \mathbb{O}_{Y_d} \rightarrow \mathbb{O}_{F_1+\dots+F_k} \rightarrow 0.$$

(Notice that the map $H^1(\mathbb{O}_{Y_d}) \rightarrow H^1(\mathbb{O}_{F_1+\dots+F_k})$ is 0, since the curves F_i are contracted by the Albanese map). \square

Remark 2.5.2. Due to the method of proof, all the surfaces constructed in the proof of Proposition 2.5.1 have an irrational pencil. The examples in Section 2.4 show that, for instance, 4, 6, $\frac{48}{7}$, 8 are accumulation points for the ratio K^2/χ of irregular surfaces without irrational pencils. We have no further information on the distribution of the ratios K^2/χ for these surfaces.

3. The Castelnuovo–de Franchis inequality

Let S be an irregular minimal surface of general type. Set $V := H^0(\Omega_S^1)$ and denote by $w : \wedge^2 V \rightarrow H^0(\omega_S)$ the natural map. If $f : S \rightarrow B$ is a pencil of genus $b \geq 2$, then $f^*H^0(\Omega_B^1)$ is a subspace of V such that $\wedge^2 f^*H^0(\Omega_B^1)$ is contained in $\ker w$ (Section 2.1). Conversely if $p_g = h^0(\omega_S) < 2q - 3$, the intersection of $\ker w$ with the cone of decomposable elements is nonzero and by 2.1.1, S has a pencil of genus $b \geq 2$. Thus:

Theorem 3.0.3 (Castelnuovo–de Franchis inequality). *Let S be an irregular surface of general type having no irrational pencil of genus $b \geq 2$. Then $p_g \geq 2q - 3$.*

In fact, using this theorem and positivity properties of the relative canonical bundle of a fibration (Theorem 4.1.1), Beauville showed:

Theorem 3.0.4 [Beauville 1982]. *Let S be a minimal surface of general type. Then $p_g \geq 2q - 4$ and if equality holds then S is the product of a curve of genus 2 and a curve of genus $q - 2 \geq 2$.*

So surfaces satisfying $p_g = 2q - 4$ have a particularly simple structure and it is natural to ask what are the irregular surfaces satisfying $p_g = 2q - 3$. Those having an irrational pencil have again a simple structure, as explained in the following theorem, which was proven in [Catanese et al. 1998] for $q = 3$, in [Barja et al. 2007] for $q = 4$ and for $q \geq 5$ in [Mendes Lopes and Pardini 2010].

Theorem 3.0.5. *Let S be a minimal surface of general type satisfying $p_g = 2q - 3$. If S has an irrational pencil of genus $b \geq 2$, then there are the following possibilities for S :*

- (i) $S = (C \times F)/\mathbb{Z}_2$, where C and F are genus 3 curves with a free involution ($q = 4$).
- (ii) S is the product of two curves of genus 3 ($q = 6$).

(iii) $S = (C \times F)/\mathbb{Z}_2$, where C is a curve of genus $2q - 3$ with a free action of \mathbb{Z}_2 , F is a curve of genus 2 with a \mathbb{Z}_2 -action such that F/\mathbb{Z}_2 has genus 1 and \mathbb{Z}_2 acts diagonally on $C \times F$ ($q \geq 3$).

In particular, $K_S^2 = 8\chi$.

So the main issue is to study the case when S has no irrational pencil. Various numerical restrictions on the invariants have been obtained. For instance if $q \geq 5$, we have (see [Mendes Lopes and Pardini 2010]):

- $K_S^2 \geq 7\chi(S) - 1$.
- If $K_S^2 < 8\chi(S)$, then $|K_S|$ has fixed components and the degree of the canonical map is 1 or 3.
- If $q(S) \geq 7$ and $K^2 < 8\chi(S) - 6$, then the canonical map is birational.

However, it is hard to say whether these results are sharp, since the only known example of a surface with $p_g = 2q - 3$ and no irrational pencil is the symmetric product of a general curve of genus 3. For low values of q we have:

- if $q = 3$, then S is the symmetric product of a curve of genus 3 — see Section 2.5(a);
- if $q = 4$, then $K^2 = 16, 17$ [Barja et al. 2007; Causin and Pirola 2006];
- if $q = 5$, there exists no such surface [Lopes et al. 2012].

Theorem 3.0.3 has been generalized to the case of Kähler manifolds of arbitrary dimension:

Theorem 3.0.6 [Pareschi and Popa 2009]. *Let X be a compact Kähler manifold with $\dim X = \text{Albdim } X = n$. If there is no surjective morphism $X \rightarrow Z$ with Z a normal analytic variety such that $0 < \dim Z = \text{Albdim } Z < \min\{n, q(Z)\}$, then*

$$\chi(\omega_X) \geq q(X) - n.$$

4. The slope inequality

4.1. Relative canonical class and slope of a fibration. In this section we consider a fibration (“pencil”) $f : S \rightarrow B$ with S a smooth projective surface and B a smooth curve of genus $b \geq 0$. Recall that a fibration is *smooth* if and only if all its fibers are smooth, and it is *isotrivial* if and only if all the smooth fibers of f are isomorphic, or, equivalently, if the fibers over a nonempty open set of B are isomorphic. Isotrivial fibrations are also said to have “constant moduli”.

We assume that the general fiber F of f has genus $g \geq 2$ and that f is *relatively minimal*, namely that there is no -1 -curve contained in the fibers of f . Notice that these assumptions are always satisfied if S is minimal of general type. Notice also that given a nonminimal fibration f it is always possible to pass to a minimal one by blowing down the -1 -curves in the fibers.

The *relative canonical class* is defined by $K_f := K_S - f^*K_B$. We also write ω_f for the corresponding line bundle $\mathcal{O}_S(K_f) = \omega_S \otimes f^*\omega_B^{-1}$.

K_f has the following positivity properties:

Theorem 4.1.1 (Arakelov; cf. [Beauville 1982]). *Let f a relatively minimal fibration of genus $g \geq 2$.*

- (i) K_f is nef. Hence, $K_f^2 = K_S^2 - 8(g - 1)(b - 1) \geq 0$.
- (ii) If f is not isotrivial, then:
 - (a) $K_f^2 > 0$
 - (b) $K_f C = 0$ for an irreducible curve C of S if and only if C is a -2 -curve contained in a fiber of f .

Let $f : S \rightarrow B$ be relatively minimal and let \bar{S} be the surface obtained by contracting the -2 curves contained in the fibers of f . There is an induced fibration $\bar{f} : \bar{S} \rightarrow B$ and, since \bar{S} has canonical singularities, $K_{\bar{f}}$ is the pullback of $K_{\bar{f}} := K_{\bar{S}} - \bar{f}^*K_B$. By the Nakai criterion for ampleness, Theorem 4.1.1(ii) can be restated by saying that if \bar{f} (equivalently, f) is not isotrivial then $K_{\bar{f}}$ is ample on \bar{S} .

Given a pencil f with general fiber of genus g , the push forward $f_*\omega_f$ is a rank g vector bundle on B of degree $\chi_f = \chi(S) - (b - 1)(g - 1)$. Recall that a vector bundle E on a variety X is said to be *nef* if the tautological line bundle $\mathbb{P}(E)$ is nef.

Theorem 4.1.2 [Fujita 1978] (see also [Beauville 1982]). *Let $f : S \rightarrow B$ be a relatively minimal fibration with general fiber of genus $g \geq 2$. Then:*

- (i) $f_*\omega_f$ is nef. In particular, $\mathbb{C}_{\mathbb{P}(f_*\omega_f)}(1)^g = \chi_f \geq 0$;
- (ii) $\chi_f = 0$ if and only if f is smooth and isotrivial.

From now on we assume that the relatively minimal fibration $f : S \rightarrow B$ is not isotrivial. The *slope* of f is defined as

$$\lambda(f) := \frac{K_f^2}{\chi_f} = \frac{K_S^2 - 8(b - 1)(g - 1)}{\chi(S) - (b - 1)(g - 1)}. \tag{4.1.1}$$

By Theorems 4.1.1 and 4.1.2, $\lambda(f)$ is well defined and > 0 .

Theorem 4.1.3 (Slope inequality). *Let $f : S \rightarrow B$ be a relatively minimal fibration with fibers of genus $g \geq 2$. If f is not smooth and isotrivial, then*

$$\frac{4(g - 1)}{g} \leq \lambda(f) \leq 12.$$

The inequality $\lambda(f) \leq 12$ follows from Noether’s formula $12\chi(S) = K_S^2 + c_2(S)$ and from the well-known formula for the Euler characteristic of a fibered surface [Barth et al. 1984, Proposition II.11.24]:

$$c_2(S) = 4(b - 1)(g - 1) + \sum_{P \in T} e(F_P) - e(F), \tag{4.1.2}$$

where e denotes the topological Euler characteristic, T is the set of critical values of f , F_P is the fiber of f over the point P and F is a general fiber. For any point $P \in B$ one has $e(F_P) \geq e(F)$, with equality holding only if F_P is smooth (loc. cit.). Hence, the main content of Theorem 4.1.3 is the lower bound $\lambda(f) \geq 4(g - 1)/g$.

We have seen (Proposition 2.5.1 and Remark 2.5.2) that the ratios K^2/χ for surfaces with an irrational pencil are dense in the interval $[2, 9]$. The slope inequality gives a lower bound for this ratio in terms of the genus g of the general fiber of the pencil.

Proposition 4.1.4. *Let S be a minimal surface of general type that has an irrational pencil $f : S \rightarrow B$ with general fiber of genus g . Then*

$$K_S^2 \geq \frac{4(g - 1)}{g} \chi(S) \geq 2\chi(S).$$

In particular, if $K_S^2 = 2\chi(S)$, then $g = 2$ and B has genus 1.

Proof. Assume that the pencil S is not smooth and isotrivial. If $K_S^2 \geq 8\chi(S)$ then of course the statement holds. If $K^2 < 8\chi$, then $K_S^2/\chi(S) \geq \lambda(f)$ and the statement follows by the slope inequality.

If f is smooth and isotrivial, then by [Serrano 1996, §1], S is a quotient $(C \times D)/G$ where C and D are curves of genus ≥ 2 and G is a finite group that acts freely. In particular, $K_S^2 = 8\chi(S)$ and the inequality is satisfied also in this case.

If $K_S^2 = 2\chi(S)$, then by the previous remarks f is not smooth and isotrivial. We have

$$2 = \frac{K_S^2}{\chi(S)} \geq \lambda(f) \geq \frac{4(g - 1)}{g}.$$

It follows immediately that $g = 2$ and B has genus 1. □

Further applications of the slope inequality are given in Section 5.

4.2. History and proofs. Theorem 4.1.3 was stated and proved first in the case of hyperelliptic fibrations [Persson 1981; Horikawa 1981].

The general statement was proved in [Xiao 1987a] and, independently, in [Cornalba and Harris 1988] under the extra assumption that the fibers of f are semistable curves, i.e., nodal curves. The proof by Cornalba and Harris has been

recently generalized in [Stoppino 2008] to cover the general case. Yet another proof was given in [Moriwaki 1996]. One can also find in [Ashikaga and Konno 2002] a nice account of Xiao's proof and of Moriwaki's proof. Hence it seems superfluous to include the various proofs here.

We just wish to point out that the three methods of proof are different. Xiao's proof uses the Harder–Narasimhan filtration of the vector bundle $f_*\omega_f$ and Clifford's Lemma.

The Cornalba–Harris proof uses GIT and relies on the fact that a canonically embedded curve of genus $g \geq 3$ is stable (fibrations whose general fiber is hyperelliptic are treated separately).

Moriwaki's proof is based on Bogomolov's instability theorem for vector bundles on surfaces.

4.3. Refinements and generalizations.

(a) Fibrations attaining the lower or the upper bound for the slope. The examples constructed in Section 2.5(c) show that the lower bound for $\lambda(f)$ given in Theorem 4.1.3 is sharp. By construction, the general fiber in all these examples is hyperelliptic. This is not a coincidence: in [Konno 1993] it is proven that the general fiber of a fibration attaining the minimum possible value of the slope is hyperelliptic. On the other hand, if $\lambda(f) = 12$, then by Noether's formula one has $c_2(S) = 4(g-1)(b-1)$. Hence by (4.1.2), this happens if and only if all the fibers of f are smooth.

(b) Nonhyperelliptic fibrations. Since, as explained in (a), the minimum value of the slope is attained only by hyperelliptic fibrations, it is natural to look for a better bound for nonhyperelliptic fibrations. In [Konno 1993], such a lower bound is established for $g \leq 5$. In [Konno 1999], it is shown that the inequality

$$\lambda(f) \geq \frac{6(g-1)}{g+1}$$

holds if: (1) g is odd, (2) the general fiber of f has maximal Clifford index, (3) Green's conjecture is true for curves of genus g .

Konno's result actually holds under assumption (1) and (2), since Green's conjecture has been proved for curves of odd genus and maximal Clifford index [Voisin 2005; Hirschowitz and Ramanan 1998].

The influence of the Clifford index and of the *relative irregularity* $q_f := q(S) - b$ has been studied also in [Barja and Stoppino 2008].

(c) Fibrations with general fiber of special type. Refinements of the slope inequality have been obtained also under the assumption that the general fiber has some special geometrical property.

Konno [1996b] showed that if the general fiber of f is trigonal and $g \geq 6$, then $\lambda(f) \geq 14(g-1)/(3g+1)$. Barja and Stoppino [2009] showed that the better bound $\lambda(f) \geq (5g-6)/g$ holds if $g \geq 6$ is even and the general fiber F of f has Maroni invariant 0. (This means that the intersection of all the quadrics containing the canonical image of F is a surface of minimal degree isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$. For the definition of the Maroni invariant see [Barja and Stoppino 2009, Remark 3.1], for instance).

The case in which the general fiber of f has an involution with quotient a curve of genus γ has been considered in [Barja 2001; Barja and Zucconi 2001; Cornalba and Stoppino 2008]: the most general result in this direction, proved in the last of these papers, is the inequality

$$\lambda(f) \geq \frac{4(g-1)}{g-\gamma} \quad \text{for } g > 4\gamma + 1.$$

(d) Generalizations to higher dimensions. Let $f : X \rightarrow B$ be a fibration, where X is an n -dimensional \mathbb{Q} -Gorenstein variety and B is a smooth curve. As in the case of surfaces, one can consider the relative canonical divisor $K_f := K_X - f^*K_B$ and define the slope of f as

$$\lambda(f) := \frac{K_f^n}{\deg f_*(\mathbb{C}_X(K_f))}.$$

This situation is studied in [Barja and Stoppino 2009], where some lower bounds are obtained under quite restrictive assumptions on the fibers.

The relative numerical invariants of threefolds fibered over a curve have also been studied in [Ohno 1992] and [Barja 2000].

5. The Severi inequality

5.1. History and proofs. Severi [1932] stated the inequality that bears his name:

Theorem 5.1.1 (Severi's inequality). *If S is a minimal surface of general type with $\text{Albdim}(S) = 2$, then*

$$K_S^2 \geq 4\chi(S).$$

Unfortunately, Severi's proof contained a fatal gap, as pointed out by Catanese [1983], who posed the inequality as a conjecture. More or less at the same time, Reid [1979] made the following conjecture, which for irregular surfaces is a consequence of Theorem 5.1.1 (compare Proposition 5.3.3):

Conjecture 5.1.2 (Reid). *If S is a minimal surface of general type such that $K_S^2 < 4\chi(S)$ then either $\pi_1^{\text{alg}}(S)$ is finite or there exists a finite étale cover $S' \rightarrow S$ and a pencil $f : S' \rightarrow B$ such that the induced surjective map on the algebraic fundamental groups has finite kernel.*

(Recall that for a complex variety X the algebraic fundamental group $\pi_1^{\text{alg}}(X)$ is the profinite completion of the topological fundamental group $\pi_1(X)$).

Motivated by this conjecture, Xiao wrote the foundational paper [Xiao 1987a] on surfaces fibred over a curve, in which he proved both Severi’s inequality in the special case of a surface with an irrational pencil and the slope inequality (Section 4).

Building on the results of [Xiao 1987a], Severi’s conjecture was proven by Konno ([Konno 1996a]) for even surfaces, namely surfaces such that the canonical class is divisible by 2 in the Picard group. At the end of the 1990’s, the conjecture was almost solved by Manetti ([Manetti 2003]), who proved it under the additional assumption that the surface have ample canonical bundle. Finally, the inequality was proven in [Pardini 2005].

The proof given in [Pardini 2005] and the proof given in [Manetti 2003] for K ample are completely different. We sketch briefly both proofs:

(a) Proof for K ample [Manetti 2003]: Let $\pi : \mathbb{P}(\Omega_S^1) \rightarrow S$ be the projection and let L be the hyperplane class of $\mathbb{P}(\Omega_S^1)$. Assume for simplicity that $H^0(\Omega_S^1)$ has no base curve. Then two general elements $L_1, L_2 \in |L|$ intersect properly, hence L^2 is represented by the effective 1-cycle $L_1 \cap L_2$. One has

$$L^2(L + \pi^*K_S) = 3(K_S^2 - 4\chi(S)). \tag{5.1.1}$$

If $L + \pi^*K_S$ is nef, then Theorem 5.1.1 follows immediately by (5.1.1). However, this is not the case in general, and one is forced to analyze the cycle $L_1 \cap L_2$ more closely. One can write

$$L_1 \cap L_2 = V + \Gamma_0 + \Gamma_1 + \Gamma_2,$$

where V is a sum of fibers of π , and the Γ_i are sums of sections of π . More precisely, the support of $\pi(\Gamma_0)$ is the union of the curves contracted by the Albanese map a , the support of $\pi(\Gamma_1)$ consists of the curves not contracted by a but on which the differential of a has rank 1 and $\pi(\Gamma_2)$ is supported on curves on which the differential of a is generically nonsingular. The term $\Gamma_0(L + \pi^*K_S)$ can be < 0 , but by means of a very careful analysis of the components of Γ_0 and of the multiplicities in the vertical cycle V one can show that

$$L^2(L + \pi^*K_S) = (L_1 \cap L_2)(L + \pi^*K_S) \geq 0.$$

Unfortunately this kind of analysis does not work when $\pi(\Gamma_0)$ contains -2 -curves, hence one has to assume K_S ample.

(a) Proof [Pardini 2005]: Set $A := \text{Alb}(S)$ and fix a very ample divisor D on A . For every integer d one constructs a fibered surface $f_d : Y_d \rightarrow \mathbb{P}^1$ as follows:

(1) Consider the cartesian diagram

$$\begin{array}{ccc} S_d & \xrightarrow{p_d} & S \\ a_d \downarrow & & \downarrow a \\ A & \xrightarrow{\mu_d} & A \end{array} \quad (5.1.2)$$

where μ_d denotes multiplication by d , and let $H_d := a_d^*D$. The map p_d is a finite connected étale cover, in particular S_d is minimal of general type.

(2) Choose a general pencil $\Lambda_d \subset |H_d|$ and let $f_d : Y_d \rightarrow \mathbb{P}^1$ be the relatively minimal fibration obtained by resolving the indeterminacy of the rational map $S_d \rightarrow \mathbb{P}^1$ defined by Λ_d .

The key observation of this proof is that as d goes to infinity, the intersection numbers H_d^2 and $K_{S_d}H_d$ grow slower than $K_{S_d}^2$ and $\chi(S_d)$. As a consequence, the slope of f_d converges to the ratio $K_S^2/\chi(S)$ as d goes to infinity. Since g_d goes to infinity with d , the Severi inequality can be obtained by applying the slope inequality (Theorem 4.1.3) to the fibrations f_d and taking the limit for $d \rightarrow \infty$.

5.2. Remarks, refinements and open questions.

(a) Chern numbers of surfaces with $\text{Albdim} = 2$. Severi's inequality is sharp, since double covers of an abelian surface branched on a smooth ample curve satisfy $K^2 = 4\chi$; see Section 2.5(d). Actually, in [Manetti 2003] it is proven that these are the only surfaces with K ample, $\text{Albdim} = 2$ and $K^2 = 4\chi$. Hence it is natural to conjecture that the canonical models of surfaces with $\text{Albdim} = 2$ and $K^2 = 4\chi$ are double covers of abelian surfaces branched on an ample curve with at most simple singularities.

In addition, by Proposition 2.5.1, the ratios $K_S^2/\chi(S)$ for S a minimal surface with $\text{Albdim} S = 2$ are dense in all the admissible interval $[4, 9]$.

(b) Refinements of the inequality. Assume that the differential of the Albanese map $a : S \rightarrow A$ is nonsingular outside a finite set. Then the cotangent bundle Ω_S^1 is nef and $L^3 = 2(K_S^2 - 6\chi(S)) \geq 0$. This remark suggests that one may expect an inequality stronger than Theorem 5.1.1 to hold under some assumption on a , e.g., that a is birational. A possible way of obtaining a result of this type would be to apply in the proof of [Pardini 2005] one of the refined versions of the slope inequality; see Section 4.3(b,c). Unfortunately, one has very little control on the general fiber of the fibrations $f_d : Y_d \rightarrow \mathbb{P}^1$ constructed in the proof.

Analogously, by the result of Manetti mentioned in (a), it is natural to expect that a better bound holds for surfaces with $q > 2$ [Manetti 2003, §7] for a series of conjectures. A step in this direction is the following:

Theorem 5.2.1 [Mendes Lopes and Pardini 2011]. *Let S be a smooth surface of maximal Albanese dimension and irregularity $q \geq 5$ with K_S ample. Then*

$$K_S^2 \geq 4\chi(S) + \frac{10}{3}q - 8.$$

Furthermore, if S has no irrational pencil and the Albanese map $a : S \rightarrow A$ is unramified in codimension 1, then

$$K_S^2 \geq 6\chi(S) + 2q - 8.$$

Theorem 5.2.1 is proven by using geometrical arguments to give a lower bound for the term $L\Gamma_2$ in the proof of Severi’s inequality for K ample given in [Manetti 2003]; see Section 5.1(a). This is why one needs to assume K ample. In [Mendes Lopes and Pardini 2011], it is also shown that the bounds of Theorem 5.2.1 can be sharpened if one assumes that the Albanese map or the canonical map of S is not birational.

(c) Generalizations to higher dimensions. The proof of Theorem 5.1.1 given in [Pardini 2005] (see Section 5.1) would work in arbitrary dimension n if one had a slope inequality for varieties of dimension $n - 1$. For instance, using the results of [Barja 2000], one can prove:

Theorem 5.2.2. *Let X be a smooth projective threefold such that K_X is nef and big and $\text{Albdim } X = 3$. Then:*

- (i) $K_X^3 \geq 4\chi(\omega_X)$.
- (ii) *If $\text{Alb}(X)$ is simple, then $K_X^3 \geq 9\chi(\omega_X)$.*

Proof. We may assume $\chi(\omega_X) > 0$. (Recall $\chi(\omega_X) \geq 0$ by [Green and Lazarsfeld 1987, Corollary to Theorem 1]). Consider a fibered threefold $f : Y \rightarrow B$, where B is a smooth curve, denote by F a general fiber and assume $\chi_f := \chi(\omega_X) - \chi(\omega_B)\chi(\omega_F) \neq 0$. Following [Barja 2000] we define

$$\lambda_2(f) := \frac{(K_X - f^*K_B)^3}{\chi_f}.$$

We apply the same construction as in Section 5.1(b) to get for every positive integer d a smooth fibered threefold $f_d : Y_d \rightarrow \mathbb{P}^1$ such that $\lambda_2(f_d)$ is defined for $d \gg 0$ and converges to $\frac{K_X^3}{\chi(\omega_X)}$ for $d \rightarrow \infty$. Statement (i) now follows by applying [Barja 2000, Theorem 3.1(i)] to f_d and taking the limit for $d \rightarrow \infty$.

Statement (ii) requires a little more care. The threefold Y_d is the blow up along a smooth curve of an étale cover $Z_d \rightarrow X$. Since $A := \text{Alb}(X)$ is simple, $V^1(X) := \{P \in \text{Pic}^0(X) | h^1(-P) > 0\}$ is a finite set by the Generic Vanishing theorem of [Green and Lazarsfeld 1987]. Then there are infinitely many values of d such that $dP \neq 0$ for every $P \in V^1(X) \setminus \{0\}$. For those values $q(Y_d) =$

$q(Z_d) = q(X)$ and $A := \text{Alb}(X)$ and $\text{Alb}(Y_d) = \text{Alb}(Z_d)$ are isogenous; in particular $\text{Alb}(Z_d)$ is simple. By construction, the general fiber F_d of f_d is a surface of maximal Albanese dimension. In addition, since F_d is isomorphic to an element of the nef and big linear system $|H_d|$ (notation as in Section 5.1(b)), it follows that $q(F_d) = q(X)$ and $\text{Alb}(F_d)$ is isogenous to the simple abelian variety $\text{Alb}(Y_d)$. Hence F_d has no irrational pencil and we get statement (ii) by applying [Barja 2000, Theorem 3.1(ii)] and taking the limit for $d \rightarrow \infty$. \square

Remark 5.2.3. In order to keep the proof of Theorem 5.2.2 simple, we have made stronger assumptions than necessary: for instance one can assume that X has terminal singularities and, with some more work, one can eliminate the assumption that $\text{Alb}(X)$ is simple in (ii).

5.3. Applications. We use the following result due to Xiao Gang:

Proposition 5.3.1 [Xiao 1987a]. *Let $f : S \rightarrow B$ be a relatively minimal fibration with fibers of genus $g \geq 2$. If $\lambda(f) < 4$ and f has no multiple fibers, then there is an exact sequence*

$$1 \rightarrow N \rightarrow \pi_1^{\text{alg}}(S) \rightarrow \pi_1^{\text{alg}}(B) \rightarrow 1,$$

where $|N| \leq 2$.

We also need the following consequence of Severi’s inequality and of the slope inequality.

Lemma 5.3.2. *Let S be a minimal regular surface of general type S with $K_S^2 < 4\chi(S)$ that has an irregular étale cover $S' \rightarrow S$. Then S has a pencil $f : S \rightarrow \mathbb{P}^1$ such that*

- (i) *f has multiple fibers $m_1 F_1, \dots, m_k F_k$ with $\sum_j \frac{m_j - 1}{m_j} \geq 2$;*
- (ii) *the general fiber of f has genus $g \leq 1 + \frac{K_S^2}{4\chi(S) - K_S^2}$.*

Proof. Up to passing to the Galois closure, we may assume that $S' \rightarrow S$ is a Galois cover with Galois group G . By Severi’s inequality (Theorem 5.1.1) the Albanese map of S' is a pencil $a' : S' \rightarrow B$, where B has genus $b > 0$. Clearly G acts on f and on B , inducing a pencil $f : S \rightarrow B/G$. Since S is regular, B/G is isomorphic to \mathbb{P}^1 . Denote by \bar{G} the quotient of G that acts effectively on B . Let $y \in B$ be a ramification point of order ν of the map $B \rightarrow B/\bar{G} = \mathbb{P}^1$ and let $H < \bar{G}$ be the stabilizer of y . The group H is cyclic of order ν and it acts freely on the fiber F_y of a' over y . It follows that the fiber of f over the image x of y is a multiple fiber of multiplicity divisible by ν . Let $x_1, \dots, x_r \in \mathbb{P}^1$ be the critical values of $B \rightarrow B/\bar{G}$, let ν_i be the ramification order of x_i and let

$m_1 F_1, \dots, m_k F_k$ be the multiple fibers of f . Then by the Hurwitz formula we have

$$\sum_j \frac{m_j - 1}{m_j} \geq \sum_i \frac{v_i - 1}{v_i} = \frac{2b - 2}{|G|} + 2 \geq 2,$$

hence (i) is proven.

To prove (ii), we observe that the fibers of f are quotients (possibly by a trivial action) of the fibers of a , hence $g \leq \gamma$, where γ is the genus of a general fiber of a . In turn, by the slope inequality one has

$$\frac{K_S^2}{\chi(S)} \geq \lambda(a) \geq \frac{4(\gamma - 1)}{\gamma},$$

which gives the required bound

$$g \leq \gamma \leq 1 + \frac{K_S^2}{4\chi(S) - K_S^2}. \quad \square$$

(a) Reid’s conjecture for irregular surfaces. Severi’s inequality implies Reid’s Conjecture 5.1.2 for irregular surfaces and, more generally, surfaces that have an irregular étale cover:

Proposition 5.3.3. *Let S be a minimal surface of general type with $K_S^2 < 4\chi(S)$. If S has an irregular étale cover, then there exists a finite étale cover $S' \rightarrow S$ and a pencil $f : S' \rightarrow B$ that induces an exact sequence*

$$0 \rightarrow N \rightarrow \pi_1^{\text{alg}}(S') \rightarrow \pi_1^{\text{alg}}(B) \rightarrow 0,$$

where $|N| \leq 2$.

Proof. Let $X \rightarrow S$ be an irregular étale cover. By Theorem 5.1.1 the Albanese map of X is a pencil $a : X \rightarrow C$ and, if $S' \rightarrow X$ is étale, then the Albanese pencil of S' is obtained by pulling back the Albanese pencil of X and taking the Stein factorization. So, up to taking a suitable base change $B \rightarrow C$ and normalizing, we can pass to an étale cover $S' \rightarrow S$ such that the Albanese pencil $a' : S' \rightarrow B$ has no multiple fiber. The statement now follows by applying Proposition 5.3.1 to a' . □

Remark 5.3.4. By Proposition 5.3.3, to prove Reid’s conjecture it is enough to show:

If S is a surface with $K_S^2 < 4\chi(S)$ that has no irregular étale cover, then $\pi_1^{\text{alg}}(S)$ is finite.

This is known to be true for $K^2 < 3\chi$; see [Mendes Lopes and Pardini 2007] and references therein. In the same reference and in [Ciliberto et al. 2007] the following sharp bound on the order of $\pi_1^{\text{alg}}(S)$ is given:

If $K_S^2 < 3\chi(S)$ and S has no irregular étale cover, then $|\pi_1^{\text{alg}}(S)| \leq 9$. Furthermore, if $|\pi_1^{\text{alg}}(S)| = 8$ or 9 then $K_S^2 = 2$ and $p_g(S) = 0$ (namely, S is a Campedelli surface).

Surfaces with $K^2 = 2$, $p_g = 0$ and $|\pi_1^{\text{alg}}| = 8$ or 9 are classified in [Mendes Lopes et al. 2009] and [Lopes and Pardini 2008], respectively.

However, all the above mentioned results make essential use of Castelnuovo's inequality $K^2 \geq 3\chi - 10$ for surfaces whose canonical map is not two-to-one onto a ruled surface. Hence, different methods are needed to deal with surfaces with $3\chi \leq K^2 < 4\chi$.

(b) Surfaces with “small” K^2 . By Proposition 2.3.2, a minimal irregular surface of general type satisfies $K^2 \geq 2\chi$. Irregular surfaces on or near the line $K^2 = 2\chi$ have been studied in [Bombieri 1973; Horikawa 1981; Reid 1979; Xiao 1987a; 1987b]. As an application of Severi's inequality and of the slope inequality, we give quick proofs of some of these results.

Proposition 5.3.5. *Let S be a minimal irregular surface of general type. Then*

- (i) *If $K_S^2 = 2\chi(S)$, then $q(S) = 1$ and the fibers of the Albanese pencil $a : S \rightarrow B$ have genus 2 (Proposition 4.1.4);*
- (ii) *if $K_S^2 < \frac{8}{3}\chi(S)$, then the Albanese map is a pencil of curves of genus 2;*
- (iii) *if $K_S^2 < 3\chi(S)$, then the Albanese map is a pencil of hyperelliptic curves of genus ≤ 3 .*

Proof. By Severi's inequality, the Albanese map of S is a pencil $a : S \rightarrow B$, where B has genus b . By the slope inequality we have

$$\frac{K_S^2}{\chi(S)} \geq \lambda(a) \geq \frac{4(g-1)}{g} \geq 2, \quad (5.3.1)$$

where, as usual, g denotes the genus of a general fiber of a . If $K_S^2 = 2\chi(S)$, then all the inequalities in (5.3.1) are equalities, hence $g = 2$ and $b = 1$.

If $K_S^2 < \frac{8}{3}\chi(S)$, then (5.3.1) gives $g = 2$.

If $K_S^2 < 3\chi(S)$, then (5.3.1) gives $g \leq 3$. The general fiber of a is hyperelliptic, since otherwise $\lambda(a) \geq 3$ by [Konno 1996b, Lem. 3.1]. \square

Next we consider regular surfaces that have an irregular étale cover.

Proposition 5.3.6. *Let S be a minimal regular surface of general type. Then:*

- (i) *if $K_S^2 < \frac{8}{3}\chi(S)$, then S has no irregular étale cover;*
- (ii) *if $K_S^2 < 3\chi(S)$, S has an irregular étale cover if and only if it has a pencil of hyperelliptic curves of genus 3 with at least 4 double fibers.*

Proof. Assume that $S' \rightarrow S$ is an irregular étale cover. Then by Lemma 5.3.2, there is a pencil $f : S \rightarrow \mathbb{P}^1$ such that the general fiber of f has genus at most

$$1 + \frac{K_S^2}{4\chi(S) - K_S^2}$$

and there are multiple fibers $m_1 F_1, \dots, m_k F_k$ such that $\sum_i (m_i - 1)/m_i \geq 2$. Since by the adjunction formula a pencil of curves of genus 2 has no multiple fibers, it follows $g \geq 3$ and $K_S^2 \geq \frac{8}{3}\chi(S)$. This proves (i).

If $K_S^2 < 3\chi(S)$, then $g = 3$ and the general fiber of f is hyperelliptic (compare the proof of Proposition 5.3.5). By the adjunction formula, the multiple fibers of f are double fibers, hence there are at least 4 double fibers.

Conversely, assume that $f : S \rightarrow \mathbb{P}^1$ is a pencil and that $y_1, \dots, y_4 \in \mathbb{P}^1$ are points such that the fiber of f over y_i is double. Let $B \rightarrow \mathbb{P}^1$ be the double cover branched on y_1, \dots, y_4 and let $S' \rightarrow S$ be obtained from $B \rightarrow \mathbb{P}^1$ by taking base change with f and normalizing. The map $S' \rightarrow S$ is an étale double cover and by construction S' maps onto the genus 1 curve B , hence $q(S') \geq 1$. \square

Remark 5.3.7. With some more work, it can be shown that Proposition 5.3.6(ii) also holds for $K_S^2 = 3\chi(S)$ [Mendes Lopes and Pardini 2007, Theorem 1.1].

References

- [Ashikaga and Konno 2002] T. Ashikaga and K. Konno, “Global and local properties of pencils of algebraic curves”, pp. 1–49 in *Algebraic geometry 2000* (Azumino), Adv. Stud. Pure Math. **36**, Math. Soc. Japan, Tokyo, 2002.
- [Barja 2000] M. A. Barja, “Lower bounds of the slope of fibred threefolds”, *Internat. J. Math.* **11**:4 (2000), 461–491.
- [Barja 2001] M. A. Barja, “On the slope of bielliptic fibrations”, *Proc. Amer. Math. Soc.* **129**:7 (2001), 1899–1906.
- [Barja and Stoppino 2008] M. Á. Barja and L. Stoppino, “Linear stability of projected canonical curves with applications to the slope of fibred surfaces”, *J. Math. Soc. Japan* **60**:1 (2008), 171–192.
- [Barja and Stoppino 2009] M. Á. Barja and L. Stoppino, “Slopes of trigonal fibred surfaces and of higher dimensional fibrations”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **8**:4 (2009), 647–658.
- [Barja and Zucconi 2001] M. Á. Barja and F. Zucconi, “On the slope of fibred surfaces”, *Nagoya Math. J.* **164** (2001), 103–131.
- [Barja et al. 2007] M. A. Barja, J. C. Naranjo, and G. P. Pirola, “On the topological index of irregular surfaces”, *J. Algebraic Geom.* **16**:3 (2007), 435–458.
- [Barth et al. 1984] W. Barth, C. Peters, and A. Van de Ven, *Compact complex surfaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **4**, Springer, Berlin, 1984.
- [Beauville 1982] A. Beauville, “L’inégalité $p_g \geq 2q - 4$ pour les surfaces de type général”, 1982. Appendix in [Debarre 1982].
- [Beauville 1996] A. Beauville, *Complex algebraic surfaces*, 2nd ed., London Mathematical Society Student Texts **34**, Cambridge University Press, Cambridge, 1996.

- [Bombieri 1973] E. Bombieri, “Canonical models of surfaces of general type”, *Inst. Hautes Études Sci. Publ. Math.* **42** (1973), 171–219.
- [Carnovale and Polizzi 2009] G. Carnovale and F. Polizzi, “The classification of surfaces with $p_g = q = 1$ isogenous to a product of curves”, *Adv. Geom.* **9**:2 (2009), 233–256.
- [Catanese 1981] F. Catanese, “On a class of surfaces of general type”, pp. 269–284 in *Algebraic surfaces* (Liguori, Naples, 1981), 1981.
- [Catanese 1983] F. Catanese, “Moduli of surfaces of general type”, pp. 90–112 in *Algebraic geometry—open problems* (Ravello, 1982), Lecture Notes in Math. **997**, Springer, Berlin, 1983.
- [Catanese 1991] F. Catanese, “Moduli and classification of irregular Kaehler manifolds (and algebraic varieties) with Albanese general type fibrations”, *Invent. Math.* **104**:2 (1991), 263–289.
- [Catanese and Ciliberto 1991] F. Catanese and C. Ciliberto, “Surfaces with $p_g = q = 1$ ”, pp. 49–79 in *Problems in the theory of surfaces and their classification* (Cortona, 1988), Sympos. Math. **32**, Academic Press, London, 1991.
- [Catanese and Ciliberto 1993] F. Catanese and C. Ciliberto, “Symmetric products of elliptic curves and surfaces of general type with $p_g = q = 1$ ”, *J. Algebraic Geom.* **2**:3 (1993), 389–411.
- [Catanese et al. 1998] F. Catanese, C. Ciliberto, and M. Mendes Lopes, “On the classification of irregular surfaces of general type with nonbirational bicanonical map”, *Trans. Amer. Math. Soc.* **350**:1 (1998), 275–308.
- [Causin and Pirola 2006] A. Causin and G. P. Pirola, “Hermitian matrices and cohomology of Kähler varieties”, *Manuscripta Math.* **121**:2 (2006), 157–168.
- [Chen and Hacon 2006] J. A. Chen and C. D. Hacon, “A surface of general type with $p_g = q = 2$ and $K_X^2 = 5$ ”, *Pacific J. Math.* **223**:2 (2006), 219–228.
- [Ciliberto and Mendes Lopes 2002] C. Ciliberto and M. Mendes Lopes, “On surfaces with $p_g = q = 2$ and non-birational bicanonical maps”, *Adv. Geom.* **2**:3 (2002), 281–300.
- [Ciliberto et al. 2007] C. Ciliberto, M. M. Lopes, and R. Pardini, “Surfaces with $K^2 < 3\chi$ and finite fundamental group”, *Math. Res. Lett.* **14**:6 (2007), 1069–1086.
- [Cornalba and Harris 1988] M. Cornalba and J. Harris, “Divisor classes associated to families of stable varieties, with applications to the moduli space of curves”, *Ann. Sci. École Norm. Sup. (4)* **21**:3 (1988), 455–475.
- [Cornalba and Stoppino 2008] M. Cornalba and L. Stoppino, “A sharp bound for the slope of double cover fibrations”, *Michigan Math. J.* **56**:3 (2008), 551–561.
- [Debarre 1982] O. Debarre, “Inégalités numériques pour les surfaces de type général”, *Bull. Soc. Math. France* **110**:3 (1982), 319–346.
- [Fujita 1978] T. Fujita, “On Kähler fiber spaces over curves”, *J. Math. Soc. Japan* **30**:4 (1978), 779–794.
- [Gieseker 1977] D. Gieseker, “Global moduli for surfaces of general type”, *Invent. Math.* **43**:3 (1977), 233–282.
- [Green and Lazarsfeld 1987] M. Green and R. Lazarsfeld, “Deformation theory, generic vanishing theorems, and some conjectures of Enriques, Catanese and Beauville”, *Invent. Math.* **90**:2 (1987), 389–407.
- [Hacon and Pardini 2002] C. D. Hacon and R. Pardini, “Surfaces with $p_g = q = 3$ ”, *Trans. Amer. Math. Soc.* **354**:7 (2002), 2631–2638.
- [Hirschowitz and Ramanan 1998] A. Hirschowitz and S. Ramanan, “New evidence for Green’s conjecture on syzygies of canonical curves”, *Ann. Sci. École Norm. Sup. (4)* **31**:2 (1998), 145–152.

- [Hirzebruch 1983] F. Hirzebruch, “Arrangements of lines and algebraic surfaces”, pp. 113–140 in *Arithmetic and geometry*, Progr. Math. **36**, Birkhäuser, Boston, 1983.
- [Horikawa 1977] E. Horikawa, “On algebraic surfaces with pencils of curves of genus 2”, pp. 79–90 in *Complex analysis and algebraic geometry*, Iwanami Shoten, Tokyo, 1977.
- [Horikawa 1981] E. Horikawa, “Algebraic surfaces of general type with small c_1^2 . V”, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **28**:3 (1981), 745–755.
- [Ishida 1983] M.-N. Ishida, “The irregularities of Hirzebruch’s examples of surfaces of general type with $c_1^2 = 3c_2$ ”, *Math. Ann.* **262**:3 (1983), 407–420.
- [Konno 1993] K. Konno, “Nonhyperelliptic fibrations of small genus and certain irregular canonical surfaces”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **20**:4 (1993), 575–595.
- [Konno 1996a] K. Konno, “Even canonical surfaces with small K^2 . III”, *Nagoya Math. J.* **143** (1996), 1–11.
- [Konno 1996b] K. Konno, “A lower bound of the slope of trigonal fibrations”, *Internat. J. Math.* **7**:1 (1996), 19–27.
- [Konno 1999] K. Konno, “Clifford index and the slope of fibered surfaces”, *J. Algebraic Geom.* **8**:2 (1999), 207–220.
- [Lopes and Pardini 2008] M. M. Lopes and R. Pardini, “Numerical Campedelli surfaces with fundamental group of order 9”, *J. Eur. Math. Soc. (JEMS)* **10**:2 (2008), 457–476.
- [Lopes et al. 2012] M. M. Lopes, R. Pardini, and G. P. Pirola, “On surfaces of general type with $q = 5$ ”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* (2012). To appear.
- [Manetti 2003] M. Manetti, “Surfaces of Albanese general type and the Severi conjecture”, *Math. Nachr.* **261/262** (2003), 105–122.
- [Mendes Lopes and Pardini 2007] M. Mendes Lopes and R. Pardini, “On the algebraic fundamental group of surfaces with $K^2 \leq 3\chi$ ”, *J. Differential Geom.* **77**:2 (2007), 189–199.
- [Mendes Lopes and Pardini 2010] M. Mendes Lopes and R. Pardini, “On surfaces with $p_g = 2q - 3$ ”, *Adv. Geom.* **10**:3 (2010), 549–555.
- [Mendes Lopes and Pardini 2011] M. Mendes Lopes and R. Pardini, “Severi type inequalities for irregular surfaces with ample canonical class”, *Comment. Math. Helv.* **86**:2 (2011), 401–414.
- [Mendes Lopes et al. 2009] M. Mendes Lopes, R. Pardini, and M. Reid, “Campedelli surfaces with fundamental group of order 8”, *Geom. Dedicata* **139** (2009), 49–55.
- [Mistretta and Polizzi 2010] E. Mistretta and F. Polizzi, “Standard isotrivial fibrations with $p_g = q = 1$, II”, *J. Pure Appl. Algebra* **214**:4 (2010), 344–369.
- [Miyaoka 1984] Y. Miyaoka, “The maximal number of quotient singularities on surfaces with given numerical invariants”, *Math. Ann.* **268**:2 (1984), 159–171.
- [Moriwaki 1996] A. Moriwaki, “A sharp slope inequality for general stable fibrations of curves”, *J. Reine Angew. Math.* **480** (1996), 177–195.
- [Ohno 1992] K. Ohno, “Some inequalities for minimal fibrations of surfaces of general type over curves”, *J. Math. Soc. Japan* **44**:4 (1992), 643–666.
- [Pardini 1991] R. Pardini, “Abelian covers of algebraic varieties”, *J. Reine Angew. Math.* **417** (1991), 191–213.
- [Pardini 2005] R. Pardini, “The Severi inequality $K^2 \geq 4\chi$ for surfaces of maximal Albanese dimension”, *Invent. Math.* **159**:3 (2005), 669–672.
- [Pareschi and Popa 2009] G. Pareschi and M. Popa, “Strong generic vanishing and a higher-dimensional Castelnuovo–de Franchis inequality”, *Duke Math. J.* **150**:2 (2009), 269–285.

- [Persson 1981] U. Persson, “Chern invariants of surfaces of general type”, *Compositio Math.* **43**:1 (1981), 3–58.
- [Pignatelli 2009] R. Pignatelli, “Some (big) irreducible components of the moduli space of minimal surfaces of general type with $p_g = q = 1$ and $K^2 = 4$ ”, *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl.* **20**:3 (2009), 207–226.
- [Pirola 2002] G. P. Pirola, “Surfaces with $p_g = q = 3$ ”, *Manuscripta Math.* **108**:2 (2002), 163–170.
- [Polizzi 2008] F. Polizzi, “On surfaces of general type with $p_g = q = 1$ isogenous to a product of curves”, *Comm. Algebra* **36**:6 (2008), 2023–2053.
- [Polizzi 2009] F. Polizzi, “Standard isotrivial fibrations with $p_g = q = 1$ ”, *J. Algebra* **321**:6 (2009), 1600–1631.
- [Reid 1979] M. Reid, “ π_1 for surfaces with small K^2 ”, pp. 534–544 in *Algebraic geometry* (Copenhagen, 1978), Lecture Notes in Math. **732**, Springer, Berlin, 1979.
- [Rito 2007] C. Rito, “On surfaces with $p_g = q = 1$ and non-ruled bicanonical involution”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **6**:1 (2007), 81–102.
- [Rito 2010a] C. Rito, “Involutions on surfaces with $p_g = q = 1$ ”, *Collect. Math.* **61**:1 (2010), 81–106.
- [Rito 2010b] C. Rito, “On equations of double planes with $p_g = q = 1$ ”, *Math. Comp.* **79**:270 (2010), 1091–1108.
- [Samuel 1966] P. Samuel, “Compléments à un article de Hans Grauert sur la conjecture de Mordell”, *Inst. Hautes Études Sci. Publ. Math.* **29** (1966), 55–62.
- [Serrano 1996] F. Serrano, “Isotrivial fibred surfaces”, *Ann. Mat. Pura Appl. (4)* **171** (1996), 63–81.
- [Severi 1932] F. di Severi, “La serie canonica e la teoria delle serie principali di gruppi di punti sopra una superficie algebrica”, *Comment. Math. Helv.* **4**:1 (1932), 268–326.
- [Sommese 1984] A. J. Sommese, “On the density of ratios of Chern numbers of algebraic surfaces”, *Math. Ann.* **268**:2 (1984), 207–221.
- [Stoppino 2008] L. Stoppino, “Slope inequalities for fibred surfaces via GIT”, *Osaka J. Math.* **45**:4 (2008), 1027–1041.
- [Voisin 2005] C. Voisin, “Green’s canonical syzygy conjecture for generic curves of odd genus”, *Compos. Math.* **141**:5 (2005), 1163–1190.
- [Xiao 1987a] G. Xiao, “Fibered algebraic surfaces with low slope”, *Math. Ann.* **276**:3 (1987), 449–466.
- [Xiao 1987b] G. Xiao, “Hyperelliptic surfaces of general type with $K^2 < 4\chi$ ”, *Manuscripta Math.* **57**:2 (1987), 125–148.
- [Yau 1977] S. T. Yau, “Calabi’s conjecture and some new results in algebraic geometry”, *Proc. Nat. Acad. Sci. U.S.A.* **74**:5 (1977), 1798–1799.
- [Zucconi 2003] F. Zucconi, “Surfaces with $p_g = q = 2$ and an irrational pencil”, *Canad. J. Math.* **55**:3 (2003), 649–672.

mmlopes@math.ist.utl.pt

*Departamento de Matemática, Instituto Superior Técnico,
Universidade Técnica de Lisboa, Av. Rovisco Pais,
1049-001 Lisboa, Portugal*

pardini@dm.unipi.it

*Dipartimento di Matematica Università di Pisa,
Largo B. Pontecorvo, 5, 56127 Pisa, Italy*